

Primordial black holes and scalar induced density perturbations: the effects of probability density functions*

Jing-Zhi Zhou (周敬之)¹ Yu-Ting Kuang (况宇庭)^{2,3†} Zhe Chang (常哲)^{2,3}

Xukun Zhang (张旭坤)⁴ Qing-Hua Zhu (朱庆华)⁵

¹Center for Joint Quantum Studies and Department of Physics, School of Science, Tianjin University, Tianjin 300350, China

²Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴College of Physics and Information Engineering, Shanxi Normal University, Taiyuan 030031, China

⁵Department of Physics and Chongqing Key Laboratory for Strongly Coupled Physics, Chongqing University, Chongqing 401331, China

Abstract: We investigate the second order energy density perturbation $\delta^{(2)}$ induced by small-scale Gaussian and local-type non-Gaussian primordial curvature perturbations. The relative abundance of primordial black holes (PBHs) is calculated in terms of the probability density functions (PDFs) of the total energy density perturbation $\delta_r = \delta^{(1)} + \frac{1}{2}\delta^{(2)}$. The effects of the second order density perturbation greatly reduce the upper bounds of small-scale power spectra of primordial curvature perturbations by one to two orders of magnitude. For a log-normal primordial power spectrum, its amplitude A_ζ is constrained to be approximately $A_\zeta \sim 3 \times 10^{-3}$, and for the local-type non-Gaussianity with $f_{\text{NL}} = 10$, the upper bound of A_ζ is approximately 2.5×10^{-4} .

Keywords: non-Gaussianity, primordial black hole, primordial power spectrum

DOI: 10.1088/1674-1137/ad972c **CSTR:** 32044.14.ChinesePhysicsC.49025105

I. INTRODUCTION

In inflation theory, valuable information about the early Universe is encoded in the cosmological perturbations that originated from the quantum fluctuations during inflation. The power spectrum of primordial curvature perturbations $\mathcal{P}_\zeta(k)$ is one of the most important predictions in inflation theory. The amplitude of the primordial power spectrum at different scales can be constrained by current cosmological observations. On large scales (≥ 1 Mpc), according to the current observations of the CMB and LSS [1, 2], the amplitude of the power spectrum of primordial curvature perturbations is constrained to be approximately 2×10^{-9} . However, on small scales (≤ 1 Mpc), the constraints of primordial curvature perturbations are significantly weaker than those on large scales [3]. Constraining the primordial power spectrum on small scales is an important issue in the study of the early Universe.

Two approaches are adopted to constrain or probe the primordial power spectrum on small scales: primordial black hole (PBH) abundance and scalar induced gravitational waves (SIGWs). More precisely, the large amplitudes of the small-scale primordial spectrum have at-

tracted significant attention in the past few years on account of their rich and profound phenomenology, such as the PBH [4–28], which is a candidate of DM. The abundance of PBHs, f_{pbh} , can be calculated in terms of a given primordial power spectrum of curvature perturbation, $\mathcal{P}_\zeta(k)$. The abundance of PBHs, $f_{\text{pbh}}(m_{\text{pbh}})$, has been constrained on different masses of PBHs, m_{pbh} [8, 29]; then, the corresponding small-scale primordial power spectrum can also be constrained by various observations of PBHs. Furthermore, the primordial perturbations will inevitably generate higher order perturbations, such as SIGWs [28, 30–60]. This strong correlation between primordial curvature perturbation and SIGW signals might be a promising approach to detecting a small-scale primordial power spectrum in the GW experiments, such as LISA and PTA [60–74].

In this study, we investigate the contributions of second order induced energy density perturbations, $\delta^{(2)} = \rho^{(2)}/\rho^{(0)}$, to the PBH formation. More precisely, the primordial curvature perturbation that enters the Hubble radius during the radiation-dominated (RD) era will lead to the generation of higher order scalar and energy density perturbations. Then, the relative abundance of PBHs can be studied in terms of the probability density functions

Received 14 October 2024; Accepted 25 November 2024; Published online 26 November 2024

* Supported by the National Nature Science Foundation of China (12475075, 12075249, 11690022) and the Key Research Program of the Chinese Academy of Sciences (XDPB15)

† E-mail: kuangyt@ihep.ac.cn

©2025 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd. All rights, including for text and data mining, AI training, and similar technologies, are reserved.

(PDFs) of the total energy density perturbations: $P(\delta^{(1)} + \delta^{(2)}/2)$, where $\delta^{(1)}$ denotes the contributions of the primordial curvature perturbation which have been studied for many years [9, 75]. Here, $\delta^{(2)}$ is the new contribution of the second order scalar induced energy density perturbation $\delta^{(2)}$, which was neglected in previous studies of PBHs¹⁾. Because the abundance of PBHs, $f_{\text{pbh}}(m_{\text{pbh}})$, has been constrained on different m_{pbh} , the amplitude of small-scale primordial power spectra can be constrained by the abundance of PBHs.

This paper is organized as follows. In Sec. II, we calculate the second order scalar induced density perturbation in the comoving gauge. In Sec. III, we study the impact of second-order scalar-induced density perturbations on the PDF and discuss the constraints on the PBH abundance imposed by the small-scale primordial power spectrum. Finally, we summarize our results and present some discussions in Sec. IV.

II. SECOND ORDER INDUCED ENERGY DENSITY PERTURBATION

The perturbed metric in the FLRW spacetime with the comoving gauge takes the form

$$ds^2 = a^2 \left(-(1 + 2\phi^{(1)} + \phi^{(2)}) d\eta^2 + (2\partial_i B^{(1)} + \partial_i B^{(2)}) d\eta dx^i + (1 - 2\psi^{(1)} - \psi^{(2)} \delta_{ij}) dx^i dx^j \right), \quad (1)$$

where $\phi^{(n)}$, $\psi^{(n)}$, and $B^{(n)}$ ($n = 1, 2$) are the n th-order scalar perturbations. We set the n -th order perturbation of the scalar part of the velocity, $\delta u^{(n)} = 0$, and the scalar metric perturbation, $E^{(n)} = 0$, in the comoving gauge. In the radiation dominated (RD) era, the first order scalar perturbation is given by

$$\begin{aligned} \psi(\eta, \mathbf{k}) &= \frac{2}{3} \zeta_{\mathbf{k}} T_{\psi}(k\eta), & \phi(\eta, \mathbf{k}) &= \frac{2}{3} \zeta_{\mathbf{k}} T_{\phi}(k\eta), \\ B(\eta, \mathbf{k}) &= \frac{2}{3k} \zeta_{\mathbf{k}} T_B(k\eta), \end{aligned}$$

where $\zeta_{\mathbf{k}}$ denotes the primordial curvature perturbations. The transfer functions $T_{\phi}(k\eta)$, $T_{\psi}(k\eta)$, and $T_B(k\eta)$ in the RD era are [76]

$$\begin{aligned} T_{\psi}(x) &= \frac{3}{2} \frac{\sin(x/\sqrt{3})}{x/\sqrt{3}}, \\ T_{\phi}(x) &= \frac{3}{2} \left(\frac{\sin(x/\sqrt{3})}{x/\sqrt{3}} - \cos(x/\sqrt{3}) \right), \\ T_B(x) &= \frac{3}{2x^2} \left(6x \cos(x/\sqrt{3}) + \sqrt{3} (x^2 - 6) \sin(x/\sqrt{3}) \right), \end{aligned}$$

where we define $x \equiv k\eta$. The higher order cosmological perturbations can be studied in terms of the xPand package [77, 78]. The equations of motion of second order scalar perturbations are

$$2\mathcal{H}\phi^{(2)'} + 6\mathcal{H}\psi^{(2)'} + 2\psi^{(2)''} + \frac{8}{3}\mathcal{H}\Delta B^{(2)} + \Delta B^{(2)'} + \Delta\phi^{(2)} - \frac{5}{3}\Delta\psi^{(2)} = -\mathcal{T}^{ij} S_{ij}^{(2)}, \quad (2)$$

$$\begin{aligned} -\mathcal{H}B^{(2)} - \frac{1}{2}B^{(2)'} - \frac{1}{2}\phi^{(2)} + \frac{1}{2}\psi^{(2)} \\ = -\Delta^{-1} \left(\partial^i \Delta^{-1} \partial^j - \frac{1}{2} \mathcal{T}^{ij} \right) S_{ij}^{(2)}, \end{aligned} \quad (3)$$

$$\partial^i (\mathcal{H} \partial_i \phi^{(2)} + \partial_i \psi^{(2)'} + S_i^{(2)}) = 0, \quad (4)$$

where $S_{ij}^{(2)}$ and $S_i^{(2)}$ are second order source terms

$$\begin{aligned} S_{ij}^{(2)} = & \delta_{ij} \left(-8\mathcal{H}\phi\phi' - 4\mathcal{H}\phi'\psi - 12\mathcal{H}\phi\psi' - 2\phi'\psi' - 4\phi\psi'' - \frac{16}{3}\mathcal{H}\phi\Delta B - \phi'\Delta B - \frac{5}{3}\psi'\Delta B \right. \\ & - 2\phi\Delta B' - 2\phi\Delta\phi - \frac{10}{3}\psi\Delta\psi + 2\mathcal{H}\partial_b B' \partial^b B - 2\mathcal{H}\partial_b \phi \partial^b B - \frac{2}{3}\mathcal{H}\partial_b \psi \partial^b B - 2\partial_b \psi' \partial^b B \\ & \left. - \frac{2}{3}\partial_b \phi \partial^b \phi + \frac{2}{3\mathcal{H}}\partial_b \psi' \partial^b \phi - 3\partial_b \psi \partial^b \psi + \frac{1}{3\mathcal{H}^2}\partial_b \psi' \partial^b \psi' - \frac{2}{3}\Delta B \Delta B + \frac{2}{3}\partial_c \partial_b B \partial^b \partial^c B \right) \\ & - \partial_b \partial_j B \partial^b \partial_i B - 2\mathcal{H}\partial_i \psi \partial_j B - \partial_i \psi' \partial_j B - \partial_i \psi \partial_j B' - \partial_i \psi \partial_j \phi - \frac{1}{\mathcal{H}}\partial_i \psi' \partial_j \phi - 2\mathcal{H}\partial_i B \partial_j \psi \\ & - \partial_i B' \partial_j \psi - \partial_i \phi \partial_j \psi + 3\partial_i \psi \partial_j \psi - \partial_i B \partial_j \psi' - \frac{1}{\mathcal{H}}\partial_i \phi \partial_j \psi' - \frac{1}{\mathcal{H}^2}\partial_i \psi' \partial_j \psi' + 4\mathcal{H}\phi \partial_i \partial_j B \\ & + \phi' \partial_i \partial_j B + \psi' \partial_i \partial_j B + \Delta B \partial_i \partial_j B + 2\phi \partial_i \partial_j B' + 2\phi \partial_i \partial_j \phi + 2\psi \partial_i \partial_j \psi, \end{aligned} \quad (5)$$

1) The statistics of induced higher order energy density perturbation $\delta^{(2)}$ are highly non-Gaussian, since they are generated by first order scalar perturbations.

$$\begin{aligned}
S_i^{(2)} = & \frac{1}{2\mathcal{H}^2} \partial_b \partial_i B' \partial^b B + \frac{1}{2\mathcal{H}^2} \partial_b \partial_i \phi \partial^b B - \frac{1}{2\mathcal{H}^2} \partial_b \partial_i B \partial^b \phi + 2\phi \partial_i B - \frac{1}{\mathcal{H}} \phi' \partial_i B - 4\psi \partial_i B \\
& - \frac{3}{\mathcal{H}} \psi' \partial_i B - \frac{1}{\mathcal{H}^2} \psi'' \partial_i B - \frac{4}{3\mathcal{H}} \Delta B \partial_i B - \frac{1}{2\mathcal{H}^2} \Delta B' \partial_i B - \frac{1}{2\mathcal{H}^2} \Delta \phi \partial_i B \\
& + \frac{4}{3\mathcal{H}^2} \Delta \psi \partial_i B + \frac{1}{\mathcal{H}} \phi \partial_i \phi - \frac{2}{\mathcal{H}} \psi \partial_i \phi - \frac{1}{\mathcal{H}^2} \psi' \partial_i \phi - \frac{1}{6\mathcal{H}^2} \Delta B \partial_i \phi + \frac{2}{3\mathcal{H}^3} \Delta \psi \partial_i \phi \\
& - \frac{2}{\mathcal{H}^2} \psi' \partial_i \psi - \frac{1}{\mathcal{H}^2} \phi \partial_i \psi' - \frac{4}{\mathcal{H}^2} \psi \partial_i \psi' - \frac{2}{\mathcal{H}^3} \psi' \partial_i \psi' - \frac{2}{3\mathcal{H}^3} \Delta B \partial_i \psi' + \frac{2}{3\mathcal{H}^4} \Delta \psi \partial_i \psi'. \quad (6)
\end{aligned}$$

\mathcal{T}^{ij} is defined as $\mathcal{T}^{ij} = \delta^{ij} - \partial^i \Delta^{-1} \partial^j$. The second order energy density perturbations $\delta^{(2)} = \rho^{(2)}/\rho^{(0)}$ can be calculated in terms of second order induced scalar perturbations and first order scalar perturbations,

$$\delta^{(2)} = -2\phi^{(2)} + \frac{2}{3\mathcal{H}^2} \Delta \psi^{(2)} - \frac{2}{3\mathcal{H}} \Delta B^{(2)} - \frac{2}{\mathcal{H}} \psi^{(2)'} + S_\rho^{(2)}, \quad (7)$$

where $S_\rho^{(2)}$ are given by

$$\begin{aligned}
S_\rho^{(2)} = & 8\phi^2 + \frac{8}{\mathcal{H}} \phi \psi' - \frac{8}{\mathcal{H}} \psi \psi' + \frac{2}{\mathcal{H}^2} (\psi')^2 \\
& + \frac{8}{3\mathcal{H}} \phi \Delta B - \frac{8}{3\mathcal{H}} \psi \Delta B + \frac{4}{3\mathcal{H}^2} \psi' \Delta B \\
& + \frac{16}{3\mathcal{H}^2} \psi \Delta \psi - 2\partial_b B \partial^b B + \frac{4}{3\mathcal{H}} \partial_b \psi \partial^b B \\
& - \frac{2}{3\mathcal{H}^2} \partial_b \phi \partial^b \phi - \frac{4}{3\mathcal{H}^3} \partial_b \psi' \partial^b \phi \\
& + \frac{2}{\mathcal{H}^2} \partial_b \psi \partial^b \psi - \frac{2}{3\mathcal{H}^4} \partial_b \psi' \partial^b \psi' \\
& + \frac{1}{3\mathcal{H}^2} \Delta B \Delta B - \frac{1}{3\mathcal{H}^2} \partial_c \partial_b B \partial^c \partial^b B. \quad (8)
\end{aligned}$$

III. UPPER BOUNDS ON PRIMORDIAL CURVATURE PERTURBATION

As the large-amplitude primordial curvature perturbations $\zeta_{\mathbf{k}}$ on small scales are necessary for the formation of PBHs, we consider the log-normal primordial power spectrum

$$\mathcal{P}_\zeta(k) = \frac{A_\zeta}{\sqrt{2\pi\sigma_*^2}} \exp\left(-\frac{\ln(k/k_*)^2}{2\sigma_*^2}\right). \quad (9)$$

In the limit of $\sigma_* \rightarrow 0$, $\mathcal{P}_\zeta(k)$ approaches to a monochromatic power spectrum, namely, $\mathcal{P}_\zeta(k) = A_\zeta k_* \delta(k - k_*)$.

In previous studies on PBHs, the PDF of the primordial curvature perturbation $\zeta_{\mathbf{k}}$ was often assumed to be Gaussian, resulting in a Gaussian PDF for the first order energy density perturbation $\delta^{(1)}$ [75]. In this study, we

consider the contribution of second order scalar induced energy density perturbations $\delta^{(2)}$ to the formation of PBHs. As we mentioned, the statistics of second order scalar induced energy density perturbations are highly non-Gaussian. Therefore, the PDFs of the total energy density perturbation $\delta_r = \delta^{(1)} + \frac{1}{2}\delta^{(2)}$ cannot be directly and simply obtained. We follow the method described in Ref. [58] to calculate the highly non-Gaussian PDFs of δ_r . In Ref. [58], the PDFs of the second order energy density perturbation induced by primordial gravitational waves were studied. Here, we calculate the PDFs of the total energy density perturbation $\delta_r = \delta^{(1)} + \frac{1}{2}\delta^{(2)}$ induced by primordial curvature perturbations $\zeta_{\mathbf{k}}$. The relation between δ_r and $\zeta_{\mathbf{k}}$ can be rewritten as

$$\begin{aligned}
\delta_r = & \delta^{(1)} + \frac{1}{2}\delta^{(2)} \\
= & \frac{2(dk)^3}{3(2\pi)^{3/2}} \sum_{\mathbf{k}_i} \frac{\mathcal{P}_\zeta(k_*|\mathbf{k}_i)}{A_\zeta} W(|\mathbf{k}_i|R) \zeta(\mathbf{k}_i) \delta^{(1)}(|\mathbf{k}_i|\eta) \\
& + \frac{2(dk)^6}{9(2\pi)^3} \sum_{\mathbf{k}_i, \mathbf{k}_j} \frac{\mathcal{P}_\zeta(k_*|\mathbf{k}_i)}{A_\zeta} \frac{\mathcal{P}_\zeta(k_*|\mathbf{k}_j)}{A_\zeta} W(|\mathbf{k}_i + \mathbf{k}_j|R) \\
& \times \zeta(\mathbf{k}_i) \zeta(\mathbf{k}_j) I_{\delta^{(2)}}(\mathbf{k}_i, \mathbf{k}_j, \eta), \quad (10)
\end{aligned}$$

where $W(x) = 3(\sin x - x \cos x)/x^3$ is the window function [58]. $I_{\delta^{(2)}}$ is the kernel function of $\delta^{(2)}$, and \mathcal{P}_ζ is the log-normal primordial power spectrum. Eq. (10) describes the relationship between the random variables δ_r and ζ . The PDF of δ_r (non-Gaussian distribution) can be simulated in terms of the PDF of ζ . The mass-scale relation of PBHs can be expressed as follows [58]:

$$M_{\text{PBH}} = 2.2 \times 10^{13} M_\odot \left(\frac{k}{1 \text{Mpc}^{-1}}\right)^{-2}. \quad (11)$$

To intuitively show the trend of the PDFs with respect to σ_* , we calculate the PDFs corresponding to different σ_* values near $\sigma_* = 0$. As shown in Fig. 1, the PDFs of δ_r are highly non-Gaussian and contract gradually

1) We provide the specific expression for $I_{\delta^{(2)}}$ in the Python program available on the following website: <https://github.com/MrFastThree/Delta-Python-Calculation-Program.git>.




Fig. 1. (color online) PDFs of the total energy density perturbation $\delta_r = \delta^{(1)} + \frac{1}{2}\delta^{(2)}$ for a log-normal primordial power spectrum with different σ_* , where we set $A_\zeta = 1$ and $\eta \approx 0.4$.

as σ_* increases. Therefore, relying solely on the variance, σ^2 , is insufficient to evaluate the probability of PBH formation. In this study, we calculate the PDF function corresponding to different values of η within the range $[0.2, 10]$. As in Ref. [58], we select the moment that yields the maximum number of PBHs, resulting in a final η of approximately 0.4. The relative abundance of PBHs $f_{\text{pbh}}(m_{\text{pbh}})$ is equivalent to the probability that the smoothed density field exceeds the threshold δ_c [75, 79],

$$\beta(M_{\text{PBH}}) = \int_{\delta_c} d\delta_r P(\delta_r), \quad (12)$$

where the value of the threshold δ_c is, in fact, influenced

by the characteristics of the primordial perturbation (its shape and the degree of non-Gaussianity). In reality, this value may vary significantly depending on these properties [80–84]. Here, we are primarily concerned with the impact of introducing second-order density perturbations on the PDF of the final density perturbation. It is important to emphasize that our calculation of the PDF here does not depend on the choice of the threshold δ_c . For simplicity, we have set $\delta_c = 0.4$.

By substituting the PDF $P(\delta^{(1)} + \frac{1}{2}\delta^{(2)})$ of the total energy density perturbations into Eq. (12), we obtain $\beta(A_\zeta)$ as a function of A_ζ (Fig. 2). As β has been constrained at different masses of PBHs, we use Fig. 18 in Ref. [8] to determine the upper bounds of A_ζ . As shown in Fig. 3, for a log-normal primordial power spectrum, its amplitude A_ζ is constrained to be approximately $A_\zeta \sim 3 \times 10^{-3}$. The local-type non-Gaussian primordial curvature perturbation is given by [85, 86]

$$\zeta_{\mathbf{k}}^{\text{NG}} = \zeta_{\mathbf{k}}^{\text{G}} + \frac{3}{5}f_{\text{NL}} \int \frac{d^3n}{(2\pi)^{3/2}} \zeta_{\mathbf{k}-\mathbf{n}}^{\text{G}} \zeta_{\mathbf{n}}^{\text{G}}, \quad (13)$$

where ζ^{G} is the Gaussian primordial curvature perturbation. Here, we consider the lowest order contributions of local-type non-Gaussianity, namely, $\delta_{\text{NG}}^{(1)} \sim A_\zeta f_{\text{NL}}$ and $\delta_{\text{NG}}^{(2)} \sim A_\zeta^{3/2} f_{\text{NL}}$. As shown in Fig. 2, in this case, $\beta(A_\zeta, f_{\text{NL}})$ is a function of A_ζ and f_{NL} . The upper bounds of A_ζ with different f_{NL} values are given in Fig. 3(b). For $f_{\text{NL}} = 10$, the upper bound of A_ζ is approximately 2.5×10^{-4} . When we consider the second order induced energy density perturbation, the effects of primordial non-Gaussianity will greatly reduce the upper bounds of A_ζ , even for a relat-




Fig. 2. (color online) Primordial black hole abundance β as a function of the amplitude of the primordial power spectrum A_ζ . The three colored dashed lines represent the relations between β and A_ζ for different σ_* values. The three solid lines represent $\beta(A_\zeta)$ for the local-type non-Gaussian primordial curvature perturbations $\zeta_{\mathbf{k}}^{\text{NG}}$ with different f_{NL} values. The black dashed line represents the previous results from $\delta^{(1)}$ in Ref. [75].




Fig. 3. (color online) Left panel: Upper bounds of A_ζ from the total energy density perturbation δ_r with different σ_* values. The blue dashed line represents the previous results from $\delta^{(1)}$ in Ref. [75]. Right panel: Upper bounds of A_ζ for different f_{NL} values. The dashed line represents the case of a Gaussian primordial curvature perturbation.

ively small f_{NL} .

IV. CONCLUSION

In this study, we analyzed the new effects of second order scalar induced energy density perturbation $\delta^{(2)}$ on PBH formation. We calculated PDFs and β as a function of A_ζ and σ_* in terms of the log-normal primordial power spectra, where we selected the threshold $\delta_c = 0.4$ for convenience. In order to not overproduce PBHs, the parameters of primordial power spectra were constrained. For $\sigma_* \rightarrow 0$, the amplitude of primordial power spectra was constrained to be approximately $A_\zeta \sim 3 \times 10^{-3}$. With the increase in σ_* , the upper bounds of A_ζ gradually increased. The local-type non-Gaussianity was also considered in this study. For $f_{\text{NL}} = 10$, the upper bound of A_ζ was approximately 2.5×10^{-4} . The effects of primordial non-Gaussianity greatly reduced the upper bounds of A_ζ , even for a relatively small f_{NL} . In addition, as the upper bound of A_ζ was constrained to be approximately $A_\zeta \sim 3 \times 10^{-3}$ for a Gaussian primordial curvature perturb-

ation, the energy density of third order SIGWs was always less than that of second order SIGWs.

The relative abundance of PBHs $f_{\text{pbh}}(m_{\text{pbh}})$ depends on the probability that the smoothed density field exceeds the threshold δ_c [79], namely, $\beta(M_{\text{PBH}}) = \int_{\delta_c} d\delta_r P(\delta_r)$. To investigate the impact of second order induced density perturbation $\delta^{(2)}$ on β , we first need to calculate the corresponding PDFs and the threshold δ_c . The calculation of the threshold δ_c is quite complex, as it depends on the shape of the primordial power spectrum and the specific PBH model under consideration. In this study, we treated the threshold δ_c as a fixed parameter, focusing on the impact of second-order scalar-induced density perturbations on the PDFs. The effects of higher-order density perturbations on the threshold δ_c may be studied in future work.

ACKNOWLEDGEMENT

We acknowledge the xPand package [77].

References

- [1] N. Aghanim *et al.*, *Astron. Astrophys.* **641**, C4 (2020), arXiv: 1807.06209[astro-ph.CO]
- [2] E. Abdalla *et al.*, *JHEAp* **34**, 49 (2022), arXiv: 2203.06142[astro-ph.CO]
- [3] T. Bringmann, P. Scott, and Y. Akrami, *Phys. Rev. D* **85**, 125027 (2012), arXiv: 1110.2484[astro-ph.CO]
- [4] M. Sasaki, T. Suyama, T. Tanaka *et al.*, *Class. Quant. Grav.* **35**, 063001 (2018), arXiv: 1801.05235[astro-ph.CO]
- [5] B. J. Carr, *Astrophys. J.* **201**, 1 (1975)
- [6] B. J. Carr and J. E. Lidsey, *Phys. Rev. D* **48**, 543 (1993)
- [7] B. Carr, F. Kuhnel, and M. Sandstad, *Phys. Rev. D* **94**, 083504 (2016), arXiv: 1607.06077[astro-ph.CO]
- [8] B. Carr, K. Kohri, Y. Sendouda *et al.*, *Rept. Prog. Phys.* **84**, 116902 (2021), arXiv: 2002.12778[astro-ph.CO]
- [9] B. Carr and F. Kuhnel, *Ann. Rev. Nucl. Part. Sci.* **70**, 355 (2020), arXiv: 2006.02838[astro-ph.CO]
- [10] J. Garcia-Bellido and E. Ruiz Morales, *Phys. Dark Univ.* **18**, 47 (2017), arXiv: 1702.03901[astro-ph.CO]
- [11] P. Ivanov, P. Naselsky, and I. Novikov, *Phys. Rev. D* **50**, 7173 (1994)
- [12] V. De Luca, G. Franciolini, and A. Riotto, *Phys. Rev. Lett.* **126**, 041303 (2021), arXiv: 2009.08268[astro-ph.CO]
- [13] V. De Luca, G. Franciolini, P. Pani *et al.*, *JCAP* **06**, 044 (2020), arXiv: 2005.05641[astro-ph.CO]
- [14] V. Vaskonen and H. Veermäe, *Phys. Rev. Lett.* **126**, 051303 (2021), arXiv: 2009.07832[astro-ph.CO]
- [15] M. Braglia, D. K. Hazra, F. Finelli *et al.*, *JCAP* **08**, 001 (2020), arXiv: 2005.02895[astro-ph.CO]

- [16] N. Bartolo, V. De Luca, G. Franciolini *et al.*, *Phys. Rev. Lett.* **122**, 211301 (2019), arXiv: 1810.12218[astro-ph.CO]
- [17] C. T. Byrnes, M. Hindmarsh, S. Young *et al.*, *JCAP* **08**, 041 (2018), arXiv: 1801.06138[astro-ph.CO]
- [18] G. Ballesteros and M. Taoso, *Phys. Rev. D* **97**, 023501 (2018), arXiv: 1709.05565[hep-ph]
- [19] J. M. Ezquiaga, J. Garcia-Bellido, and E. Ruiz Morales, *Phys. Lett. B* **776**, 345 (2018), arXiv: 1705.04861[astro-ph.CO]
- [20] K. Inomata, M. Kawasaki, K. Mukaida *et al.*, *Phys. Rev. D* **95**, 123510 (2017), arXiv: 1611.06130[astro-ph.CO]
- [21] M. Kawasaki, A. Kusenko, Y. Tada *et al.*, *Phys. Rev. D* **94**, 083523 (2016), arXiv: 1606.07631[astro-ph.CO]
- [22] S. Young and C. T. Byrnes, *JCAP* **08**, 052 (2013), arXiv: 1307.4995[astro-ph.CO]
- [23] A. D. Gow, C. T. Byrnes, P. S. Cole *et al.*, *JCAP* **02**, 002 (2021), arXiv: 2008.03289[astro-ph.CO]
- [24] S. Wang, K. Kohri, and V. Vardanyan, (2021), arXiv: 2107.01935[gr-qc]
- [25] M. Y. Khlopov, *Res. Astron. Astrophys.* **10**, 495 (2010), arXiv: 0801.0116[astro-ph]
- [26] K. M. Belotsky, V. I. Dokuchaev, Y. N. Eroshenko *et al.*, *Eur. Phys. J. C* **79**, 246 (2019), arXiv: 1807.06590[astro-ph.CO]
- [27] S. Wang, Y. F. Wang, Q.-G. Huang *et al.*, *Phys. Rev. Lett.* **120**, 191102 (2018), arXiv: 1610.08725[astro-ph.CO]
- [28] S. Wang, T. Terada, and K. Kohri, *Phys. Rev. D* **99**, 103531 (2019), arXiv: 1903.05924[astro-ph.CO]
- [29] B. J. Carr, K. Kohri, Y. Sendouda *et al.*, *Phys. Rev. D* **81**, 104019 (2010), arXiv: 0912.5297[astro-ph.CO]
- [30] G. Domènech, *Universe* **7**, 398 (2021), arXiv: 2109.01398[gr-qc]
- [31] S. Mollerach, D. Harari, and S. Matarrese, *Phys. Rev. D* **69**, 063002 (2004), arXiv: astro-ph/0310711
- [32] K. N. Ananda, C. Clarkson, and D. Wands, *Phys. Rev. D* **75**, 123518 (2007), arXiv: gr-qc/0612013
- [33] D. Baumann, P. J. Steinhardt, K. Takahashi *et al.*, *Phys. Rev. D* **76**, 084019 (2007), arXiv: hep-th/0703290
- [34] K. Kohri and T. Terada, *Phys. Rev. D* **97**, 123532 (2018), arXiv: 1804.08577[gr-qc]
- [35] J. Z. Zhou, X. Zhang, Q. H. Zhu *et al.*, *JCAP* **05**, 013 (2022), arXiv: 2106.01641[astro-ph.CO]
- [36] Z. Chang, Y. T. Kuang, X. Zhang *et al.*, *Chin. Phys. C* **47**, 055104 (2023), arXiv: 2209.12404[astro-ph.CO]
- [37] R. Saito and J. Yokoyama, *Phys. Rev. Lett.* **102**, 161101 (2009), arXiv: 0812.4339[astro-ph]
- [38] K. Inomata and T. Nakama, *Phys. Rev. D* **99**, 043511 (2019), arXiv: 1812.00674[astro-ph.CO]
- [39] C. T. Byrnes, P. S. Cole, and S. P. Patil, *JCAP* **06**, 028 (2019), arXiv: 1811.11158[astro-ph.CO]
- [40] J. Garcia-Bellido, M. Peloso, and C. Unal, *JCAP* **09**, 013 (2017), arXiv: 1707.02441[astro-ph.CO]
- [41] T. Nakama, J. Silk, and M. Kamionkowski, *Phys. Rev. D* **95**, 043511 (2017), arXiv: 1612.06264[astro-ph.CO]
- [42] E. Bugaev and P. Klimai, *Phys. Rev. D* **83**, 083521 (2011), arXiv: 1012.4697[astro-ph.CO]
- [43] R. Saito and J. Yokoyama, *Prog. Theor. Phys.* **123**, 867 (2010), arXiv: 0912.5317[astro-ph.CO]
- [44] E. Bugaev and P. Klimai, *Phys. Rev. D* **81**, 023517 (2010), arXiv: 0908.0664[astro-ph.CO]
- [45] K. Inomata, M. Kawasaki, K. Mukaida *et al.*, *Phys. Rev. D* **101**, 123533 (2020), arXiv: 2003.10455[astro-ph.CO]
- [46] G. Ballesteros, J. Rey, M. Taoso *et al.*, *JCAP* **07**, 025 (2020), arXiv: 2001.08220[astro-ph.CO]
- [47] J. Lin, Q. Gao, Y. Gong *et al.*, *Phys. Rev. D* **101**, 103515 (2020), arXiv: 2001.05909[gr-qc]
- [48] Z.-C. Chen, C. Yuan, and Q.-G. Huang, *Phys. Rev. Lett.* **124**, 251101 (2020), arXiv: 1910.12239[astro-ph.CO]
- [49] R.-G. Cai, S. Pi, S.-J. Wang *et al.*, *JCAP* **10**, 059 (2019), arXiv: 1907.06372[astro-ph.CO]
- [50] Y.-F. Cai, C. Chen, X. Tong *et al.*, *Phys. Rev. D* **100**, 043518 (2019), arXiv: 1902.08187[astro-ph.CO]
- [51] K. Ando, K. Inomata, and M. Kawasaki, *Phys. Rev. D* **97**, 103528 (2018), arXiv: 1802.06393[astro-ph.CO]
- [52] H. Di and Y. Gong, *JCAP* **07**, 007 (2018), arXiv: 1707.09578[astro-ph.CO]
- [53] Z. Chang, X. Zhang, and J. Z. Zhou, (2022), arXiv: 2207.01231[astro-ph.CO]
- [54] Z. Chang, X. Zhang, and J. Z. Zhou, (2022), arXiv: 2209.07693[astro-ph.CO]
- [55] N. Orlofsky, A. Pierce, and J. D. Wells, *Phys. Rev. D* **95**, 063518 (2017), arXiv: 1612.05279[astro-ph.CO]
- [56] Q. Gao, *Sci. China Phys. Mech. Astron.* **64**, 280411 (2021), arXiv: 2102.07369[gr-qc]
- [57] T. Nakama and T. Suyama, *Phys. Rev. D* **92**, 121304 (2015), arXiv: 1506.05228[gr-qc]
- [58] T. Nakama and T. Suyama, *Phys. Rev. D* **94**, 043507 (2016), arXiv: 1605.04482[gr-qc]
- [59] Z. Changa, K. Y. Tinga, X. Zhang *et al.*, (2022), arXiv: 2211.11948[astro-ph.CO]
- [60] Z. C. Zhao and S. Wang, *Universe* **9**, 157 (2023), arXiv: 2211.09450[astro-ph.CO]
- [61] A. Afzal *et al.*, *Astrophys. J. Lett.* **951**, L11 (2023), arXiv: 2306.16219[astro-ph.HE]
- [62] S. Wang, Z.-C. Zhao, J. P. Li *et al.*, (2023), arXiv: 2307.00572[astro-ph.CO]
- [63] Y. C. Bi, Y. M. Wu, Z. C. Chen *et al.*, (2023), arXiv: 2307.00722[astro-ph.CO]
- [64] C. Yuan, Z. C. Chen, and Q.-G. Huang, *Phys. Rev. D* **100**, 081301 (2019), arXiv: 1906.11549[astro-ph.CO]
- [65] E. Thrane and J. D. Romano, *Phys. Rev. D* **88**, 124032 (2013), arXiv: 1310.5300[astro-ph.IM]
- [66] T. Robson, N. J. Cornish, and C. Liu, *Class. Quant. Grav.* **36**, 105011 (2019), arXiv: 1803.01944[astro-ph.HE]
- [67] K. Kuroda, W. T. Ni, and W.-P. Pan, *Int. J. Mod. Phys. D* **24**, 1530031 (2015), arXiv: 1511.00231[gr-qc]
- [68] X. Siemens, J. Ellis, F. Jenet *et al.*, *Class. Quant. Grav.* **30**, 224015 (2013), arXiv: 1305.3196[astro-ph.IM]
- [69] J. Z. Zhou, Y. T. Kuang, D. Wu *et al.*, (2024), arXiv: 2409.07702[gr-qc]
- [70] Z. Chang, Y. T. Kuang, D. Wu *et al.*, *JCAP* **2024**, 044 (2024), arXiv: 2312.14409[astro-ph.CO]
- [71] Z. Chang, Y. T. Kuang, D. Wu *et al.*, *Phys. Rev. D* **109**, L041303 (2024), arXiv: 2311.05102[astro-ph.CO]
- [72] J. Ellis, M. Fairbairn, G. Franciolini *et al.*, *Phys. Rev. D* **109**, 023522 (2024), arXiv: 2308.08546[astro-ph.CO]
- [73] G. Franciolini, A. Iovino, Junior., V. Vaskonen *et al.*, *Phys. Rev. Lett.* **131**, 201401 (2023), arXiv: 2306.17149[astro-ph.CO]
- [74] G. Domènech, S. Pi, A. Wang, and J. Wang, *JCAP* **08**, 054 (2024), arXiv: 2402.18965[astro-ph.CO]
- [75] G. Sato-Polito, E. D. Kovetz, and M. Kamionkowski, *Phys. Rev. D* **100**, 063521 (2019), arXiv: 1904.10971[astro-ph.CO]
- [76] Y. Lu, A. Ali, Y. Gong *et al.*, *Phys. Rev. D* **102**, 083503 (2020), arXiv: 2006.03450[gr-qc]

- [77] C. Pitrou, X. Roy, and O. Umeh, *Class. Quant. Grav.* **30**, 165002 (2013), arXiv: 1302.6174[astro-ph.CO]
- [78] K. Tomikawa and T. Kobayashi, *Phys. Rev. D* **101**, 083529 (2020), arXiv: 1910.01880[gr-qc]
- [79] W. H. Press and P. Schechter, *Astrophys. J.* **187**, 425 (1974)
- [80] I. Musco, *Phys. Rev. D* **100**, 123524 (2019), arXiv: 1809.02127[gr-qc]
- [81] I. Musco, V. De Luca, G. Franciolini *et al.*, *Phys. Rev. D* **103**, 063538 (2021), arXiv: 2011.03014[astro-ph.CO]
- [82] F. Noble Chamings, A. Avgoustidis, E. J. Copeland *et al.*, *Phys. Rev. D* **100**, 043525 (2019), arXiv: 1904.00884[astro-ph.CO]
- [83] V. De Luca, G. Franciolini, A. Kehagias *et al.*, *JCAP* **07**, 048 (2019), arXiv: 1904.00970[astro-ph.CO]
- [84] V. De Luca, A. Kehagias, and A. Riotto, *Phys. Rev. D* **108**, 063531 (2023), arXiv: 2307.13633[astro-ph.CO]
- [85] R.-g. Cai, S. Pi, and M. Sasaki, *Phys. Rev. Lett.* **122**, 201101 (2019), arXiv: 1810.11000[astro-ph.CO]
- [86] J. P. Li, S. Wang, Z. C. Zhao *et al.*, (2023), arXiv: 2309.07792[astro-ph.CO]