

Black holes in general relativity coupled with nonlinear electrodynamics surrounded by perfect fluid dark matter: Thermodynamics, particle motion and black hole shadow

Bekzod Rahmatov^{1,2†} Muhammad Zahid^{3‡} Saeed Ullah Khan^{4§} Javlon Rayimbaev^{5,6¶} Inomjon Ibragimov^{7#}
Zukhriddin Yuldoshev^{8¶} Adilbek Dauletov^{9^b} Sokhibjan Muminov^{10^a}

¹Samarkand State University, 140104, University blv.15, Samarkand, Uzbekistan

²Shahrisabz State Pedagogical Institute, Shahrisabz Str. 10, Shahrisabz 181301, Uzbekistan

³School of Science, Harbin Institute of Technology, ShenZhen, 518055, China

⁴School of Mathematical Sciences, Shenzhen University, Shenzhen 518060, China

⁵New Uzbekistan University, Movarounnahr Street 1, Tashkent 100007, Uzbekistan

⁶Urgench State University, Kh. Alimjan Str. 14, Urgench 221100, Uzbekistan

⁷Kimyo International University in Tashkent, Shota Rustaveli street 156, Tashkent 100121, Uzbekistan

⁸Tashkent State Technical University, Tashkent 100095, Uzbekistan

⁹Alfraganus University, Yukori Karakamish Street 2a, Tashkent 100190, Uzbekistan

¹⁰Mamun University, Bolkhovuz Street 2, Khiva 220900, Uzbekistan

Abstract: This study explores black holes in General Relativity (GR) coupled with nonlinear electrodynamics (NED) in the presence of perfect fluid dark matter (PFDM). We derive a singular black hole solution and investigate its thermodynamic properties, including black hole temperature, entropy, and specific heat capacity of the black hole spacetime. The analysis of energy conditions reveals deviations from standard GR, with PFDM affecting the weak and strong energy conditions. The study further examines the impact of NED and PFDM on the innermost stable circular orbit (ISCO), demonstrating that PFDM shifts the ISCO radius and that combined effects of NED and PFDM field parameters sufficiently influence orbital stability. Our analysis of the black hole shadow reveals that PFDM increases the shadow radius while a higher charge reduces it, leading to modifications in potential astrophysical observables. The thermodynamic behavior of the black hole exhibits phase transitions marked by changes in heat capacity, indicating possible stability regimes. Moreover, we also derive equations for black hole shadow size and study the spacetime effects on the shadow. These results provide a framework for testing alternative gravity theories and understanding the role of exotic matter in strong gravitational fields. Finally, we compare the constraints on NED and PFDM field parameters derived from our black hole model with the Event Horizon Telescope (EHT) observations of M87* and Sgr A*, providing observational limits on deviations from General Relativity.

Keywords: Singular Black Holes, PFDM field, Black hole shadow, Black hole thermodynamics, GR coupling to NED field

DOI: **CSTR:**

I. INTRODUCTION

Black holes are among the most challenging and still debated aspects of Einstein's theory of relativity. Regular or non-singular black holes have been predicted using certain energy-momentum tensors, resulting in secondary consequences. Recent research suggests that violating the

strong energy condition is a crucial characteristic of regular black holes. The hunt for regular black holes began with Sakharov and Gliner [1, 2], who hypothesized that using an environment with a de Sitter metric rather than a vacuum may prevent the occurrence of singularities. Gurevich and Starobinsky [3, 4] postulated a complex in-

Received 14 February 2025; Accepted 18 March 2025

[†] E-mail: rahmatovbekzod@samdu.uz

[‡] E-mail: zahid.m0011@gmail.com

[§] E-mail: saeedkhan.u@gmail.com

[¶] E-mail: javlon@astrin.uz

[¶] E-mail: i.ibragimov@kiut.uz

[¶] E-mail: yuldoshevzuxriddin68@gmail.com

^b E-mail: a.dauletov@afu.uz

^a E-mail: sokhibjan.muminov@mamunedu.uz

©2025 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd. All rights, including for text and data mining, AI training, and similar technologies, are reserved.

teraction between gravitational and electromagnetic fields.

Nonlinear electrodynamics (NED) can generate fields that may be coupled to potential gravitational sources, especially in the neighborhood of gigantic entities like black holes. Under these circumstances, the gravitational field functions as a generator of electromagnetic fields, intensifying them and affecting the curvature of spacetime [5]. The motion of charged particles in their vicinity may be explained using a NED. Therefore, it is essential to study a black hole in General Relativity (GR) connected to NED [6, 7].

According to recent cosmological investigations, the mainstream model of cosmology claims that dark energy (DE) accounts for almost 68.3% of our cosmos, while dark matter (DM) accounts for 26.8%. The baryon matter makes up the remaining 4.9% [8]. We infer the presence of DM in the visible cosmos from indirect evidence. As we go farther from the galactic center, its prevalence increases. These assumptions are concluded from the analysis of galaxy rotation [9].

Consequently, there is a need to examine black hole solutions encompassed by DE or DM, a topic that has recently received significant interest from researchers. Kiselev, for the first, investigated the Schwarzschild black hole surrounded by quintessential energy [10], and this study was later expanded to include the Kerr-like black hole in [11]. The perfect fluid dark matter (PFDM) model is a phenomenological framework containing a non-zero DM component represented by a logarithmic term $\alpha \ln(r/r_q)$. Rehman et al. [12] constructed a PFDM model and examined the energy density, pressure, and equation of the DM. Subsequently, several researchers engaged in the investigation of black holes in the presence of quintessence; for further information, see [13–22].

Li and Yang [23] provide an alternative black hole solution consisting of the term $(r_q/r) \ln(r/r_q)$. They do this by emphasizing that the DM halo, represented by a phantom scalar field, is composed of heavy particles with weak interactions with the equation of state. Since the phantom scalar field plays no role in the cosmos, thus this paradigm classifies background matter dispersion as DM. The contents of DE, such as a quintessence field or a cosmological constant, significantly impact the geometry of spacetime, particularly in the context of black holes. Therefore, in light of their approach, various astrophysical black hole characteristics have been explored in different DE and DM scenarios [24–33].

Black holes are famous for their strong gravitational attractions and thermodynamics. Bardeen et al. [34] introduce the four laws of black hole thermodynamics. In physics, thermodynamics is helpful due to its inherent features. This lets us study a system's macroscopic physics without knowing its details. Hawking showed that

black holes produce semiclassical thermal radiation like pure black bodies [35]. Schwarzschild black holes have a negative heat capacity and are unstable due to thermal fluctuations since their Hawking temperatures are inversely proportional to mass. These characteristics attracted many researchers to explore black hole thermodynamics for better understanding [36–40].

The dynamics of charged and neutral test particles near black holes are essential topics in relativistic astrophysics. They help us to comprehend the powerful gravitational field that black holes possess. Therefore, in this article, we aim to examine the model or solution of gravity theories by studying the particle motion around compact objects. In particular, specific X-ray sources are thought to be related to the accretion disc around astrophysical black holes. Measurements of the disc's inner edge radii may provide limits on the parameters of alternative and modified theories of gravity [41–56]. The process of capturing massive and massless particles by parameterized black holes has been extensively investigated in [57–59]. Analysis of the orbital and epicyclic frequencies in axially symmetric and stationary spacetime is studied in Refs. [60–63]. The above studies strongly encourage investigating the motion of test particles around black holes. Nonlinear electrodynamics, in conjunction with general relativistic solutions, explain this motion.

Recent studies, such as [64–66], have explored black hole shadows in NED, highlighting its effects on spacetime geometry and astrophysical observables. Inspired by these works, we incorporate PFDM to analyze its impact together with the NED field on black hole thermodynamics, particle dynamics, and shadow properties, offering deeper insights into deviations from pure GR.

The connection between black hole thermodynamics and shadow properties, as explored in [67], shows that phase transitions influence shadow size and stability. This work extends these analyses by incorporating PFDM and NED, demonstrating their combined effects on thermodynamics and shadow observables.

Syngé first explored the shadow cast by a Schwarzschild black hole [68] and later by Luminet [69]. Syngé derived a formula for determining the angular radius of the shadow. The first study on the appearance of the Kerr black hole's shadow was conducted by Bardeen [70], with the results also included in Chandrasekhar's book [71] and in [72]. Unlike the Schwarzschild case, the shadow of a Kerr black hole is not perfectly circular.

Hioki and Maeda [73] analyzed the shadow of both a Kerr black hole and a Kerr naked singularity by introducing two observables. More recently, some of the authors of this paper developed a coordinate-independent approach to characterize black hole shadows [74]. Various other black hole solutions have also been investigated in this context, including the Kerr-Newman black hole [75], the Einstein-Maxwell-Dilaton-Axion black hole [76], the

Kerr-Taub-NUT black hole [77], the rotating braneworld black hole [78], the Kaluza-Klein rotating dilaton black hole [79], the rotating non-Kerr black hole [80], and the Kerr-Newman-NUT black hole with a cosmological constant [81].

Additionally, the study of black hole shadows has been extended to higher dimensions, such as the 5D rotating Myers-Perry black hole [82]. Furthermore, a unique example of a single black hole solution in general relativity that exhibits multiple shadows was first presented in [83].

The recent Event Horizon Telescope (EHT) observations of M87* and Sgr A* provide direct tests of general relativity and constraints on black hole shadow properties [84, 85]. Modifications due to NED and PFDM have been shown to affect spacetime geometry and shadow structures [86, 87].

In this work, we analyze geodesic motion and derive black hole shadow observables in a spacetime influenced by NED and PFDM. In addition, we explore the thermodynamic behavior of the system and its observational implications and phase transitions to shadow variations [88, 89].

The primary goal of this paper is to investigate singular black holes in GR coupled to NED in the presence of PFDM. In the following Sec II, we will provide a concise overview of the spacetime geometry of black holes encircled by PFDM and coupled to NED. In Sec. IV, we will examine the features of black holes thermodynamics in the presence of PFDM. Section V is devoted to investigating neutral particle motion and their important characteristics in response to the presence of PFDM. We explore black hole shadows under the influence of PFDM in Sec. VII. Finally, in the last Sec. IX, we wrap up our investigation with concluding comments.

II. BLACK HOLES IN GR COUPLED TO NED SURROUNDED BY PFDM

In the context of black hole solutions, researchers often study black hole behavior in different theoretical frameworks or gravitational backgrounds. This could involve exploring various metrics that describe spacetime geometry around black holes, such as the Schwarzschild metric for non-rotating black holes or the Kerr metric for rotating black holes.

A. Black hole solutions with the dark fluid equation of state

The present section briefly introduces a black hole solution in GR coupled with the NED field and PFDM in a spherically symmetric spacetime with the ansatz

$$ds^2 = -e^{\nu(r)} dt^2 + e^{\Lambda(r)} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (1)$$

Nonsingular (regular) extensions of black hole spacetimes have been widely considered in GR coupled to NED [90–100]. But, here, we focus on a new black hole solution in GR coupled with NED in the presence of an exotic field called the PFDM field using the following corresponding action:

$$S = \frac{1}{16\pi} \int d^4x \sqrt{|g|} (R - \mathcal{L}_{\text{NED}} + \mathcal{L}_{\text{PFDM}}), \quad (2)$$

where, R is the Ricci scalar, g is determinant of the metric tensor $g_{\mu\nu}$, NED field Lagrangian is given in Ref.[101]

$$\mathcal{L}_{\text{NED}} = \frac{4\mu}{\alpha} \frac{(\alpha F)^{(\nu+3)/4}}{[1 + (\alpha F)^{\nu/4}]^{1+\mu/\nu}}, \quad (3)$$

with the coupling parameter α and the Faraday tensor of the electromagnetic field $F = F_{ab}F^{ab}$. Since there is no matter Lagrangian, the energy-momentum tensor in Schwarzschild coordinates is due solely to $\mathcal{L}_{\text{NED}}$ as

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = T_{\mu\nu}^{(\text{NED})} + T_{\mu\nu}^{(\text{PFDM})}, \quad (4)$$

where the perfect fluid energy-momentum tensor is

$$T_{\mu\nu}^{(\text{PF})} = (\rho + p)u_\mu u_\nu + pg_{\mu\nu}, \quad (5)$$

which, in the case of PFDM, has only the standard orthogonal basis

$$T_{\mu\nu}^{(\text{PFDM})} = \text{diag}(-\epsilon, p_r, p_\theta, p_\phi) \quad (6)$$

with

$$-\epsilon = p_r = \frac{\lambda}{8\pi r^3}, \quad p_\theta = p_\phi = -\frac{\lambda}{16\pi r^3}. \quad (7)$$

The energy-momentum tensor for the NED reads as,

$$4\pi T_{\mu\nu}^{(\text{NED})} = -2 \frac{\partial \mathcal{L}_{\text{NED}}}{\partial F} F_\mu^\beta F_{\nu\beta} + \mathcal{L}_{\text{NED}} g_{\mu\nu}. \quad (8)$$

The non-zero diagonal components of the tensor for the NED field are

$$T_0^0 = T_1^1 = -\frac{2\mu M q^\nu r^{\mu-3}}{(q^\nu + r^\nu)^{\frac{\mu+\nu}{\nu}}}, \quad (9)$$

$$T_2^2 = T_3^3 = \frac{\mu M q^\nu r^{\mu-3}}{(q^\nu + r^\nu)^{\frac{\mu}{\nu}+2}} [(v+1)r^\nu - (\mu-1)q^\nu]. \quad (10)$$

To solve this field equation, first, we find the component G_{tt} of the Einstein tensor using the following ansatz

$$e^{\nu(r)} = e^{-\Lambda(r)} = f(r) \quad (11)$$

and we get

$$G^0_0 = G^1_1 = -\frac{1}{r} \frac{df(r)}{dr} - \frac{f(r)}{r^2} + \frac{1}{r^2}, \quad (12)$$

$$G^2_2 = G^3_3 = -\frac{1}{2} \frac{d^2 f(r)}{dr^2} - \frac{1}{r} \frac{df(r)}{dr}. \quad (13)$$

For a static and spherically symmetric spacetime with line element Eq. (1) and NED source given by a magnetic charge q in which $F_{\mu\nu} = (\delta^\theta_\mu \delta^\phi_\nu - \delta^\theta_\nu \delta^\phi_\mu) q \sin \theta$.

One can have the following differential equation for the metric function $f(r)$, using Eq.(4) and taking into account Eqs. (6), (9) and (12):

$$\frac{1}{r} \frac{df(r)}{dr} + \frac{f(r)}{r^2} + \frac{1}{r^2} = \frac{2\mu M q^\nu r^{\mu-3}}{(q^\nu + r^\nu)^{\frac{\mu+\nu}{\nu}}} - \frac{\lambda}{8\pi r^3}. \quad (14)$$

The solution of Eq.(14) takes the form,

$$f(r) = \frac{-2Mr^\mu (q^\nu + r^\nu)^{-\frac{\mu}{\nu}} + \lambda \ln r + r}{r} + \frac{c_1}{r} \quad (15)$$

After simplifying Eq. (15) we can get the following form,

$$f(r) = 1 - \frac{2M}{r} \left(1 + \frac{q^\nu}{r^\nu}\right)^{-\frac{\mu}{\nu}} + \frac{\lambda \ln r}{r} + \frac{c_1}{r}. \quad (16)$$

Noting that a quantity in the logarithmic term should be dimensionless. Therefore, we chose the constant as $c_1 = -\lambda \ln |\lambda|$, and we get the following form for the metric function:

$$f(r) = 1 - \frac{2M}{r} \left(1 + \frac{q^\nu}{r^\nu}\right)^{-\frac{\mu}{\nu}} + \frac{\lambda}{r} \ln \frac{r}{|\lambda|}. \quad (17)$$

Now, we analyze the horizon properties of the obtained black hole solution graphically.

B. Horizon properties

Fig. 1 shows how the metric function $f(r)$ around the black hole changes under different parameter influences. Increasing the NED parameters reduces spacetime curvature, making the black hole more stable. A higher charge strengthens the electromagnetic field, expanding the event horizon. PFDM causes spacetime to extend and diminishes dynamic effects outside the black hole. These graphs highlight the interplay of these factors, providing insights into the structure of spacetime surrounding the black hole.

The given set of graphs in Fig. 2 represents the radius of the event horizon r_H versus different parameters (λ and q) for various combinations of the parameters ν and μ . The upper row shows r_H versus λ , and the bottom row shows r_H versus q . Each column corresponds to different values of ν and μ . The color panel on the right side of each graph shows the value of the third parameter, λ and q . The radius of the event horizon has a minimum value of a certain value of λ . The curves show all graphs. This minimum value varies according to the values of ν and μ . As the value increases, the initial value of the event horizon radius, r_H , increases on the graphs. It shows that increasing ν causes the black hole to grow larger. An increase in the value of μ also increases the value of r_H , but this effect is weaker than the impact of ν . The U-shaped curves show the radius of the minimum event horizon as λ changes. This minimum value is reached at a certain value of λ , which varies for different parameters ν and μ . The graphs show that an increase in λ first decreases and then increases the radius of the event horizon. All graphs show that with increasing q , the radius of the event horizon r_H generally decreases. This means that an increase in magnetic charge shrinks the radius of the black hole's event horizon.




Fig. 1. (color online) Radial dependence of the metric function $f(r)$ for various values of q and λ and fixed values of μ and ν .




Fig. 2. (color online) The Event horizon radius for q and λ .

C. The scalar invariants

To explore the properties of spacetime, we may start by studying the scalar invariants of geometry around a compact gravitating object. Here, we analyze the Ricci scalar, square of the Ricci tensor, and Kretschmann scalar of the spacetime metric (1) given with the lapse function given by Eq. 17.

1. Ricci scalar

The Ricci scalar, often known as the scalar curvature, is a fundamental measure of curvature in warped spacetime. The Ricci scalar measures the curvature of spacetime resulting from the presence of matter or energy, influenced by the black hole's intrinsic properties and any external fields or modifications, such as PFDM and NED field parameters. It also affects the orbits of particles in the accretion disk, such as the location of the ISCO, which sets the inner edge of the accretion disk and the orbital velocities, as well as the stability of particles in the disk. The accretion disk's emission properties depend on the disk's temperature and density profile, shaped by the spacetime curvature. In turn, changes in the Ricci scalar cause deviations in the temperature distribution, and thus, the emitted radiation will shift if the inner disk edge moves closer to or farther from the black hole. Observations of the accretion disk emissions can constrain the curvature properties and test deviations from standard GR, such as those introduced by PFDM and NED field.

Ricci scalar, the so-called scalar curvature, is one of the simplest curvature invariants of curved spacetime and is defined as $R = g^{\mu\nu} R_{\mu\nu}$, where $R_{\mu\nu}$ is the Ricci tensor. The positive and the negative values of the Ricci scalar correspond to the sunken and convex form of the spacetime, respectively. After some simple mathematics, one may quickly get the following form for the Ricci scalar

for the solution.

$$R = \frac{\left(1 + \frac{q^\nu}{r^\nu}\right)^{-\frac{\mu}{\nu}}}{r^3(q^\nu + r^\nu)^2} \left\{ -r^{2\nu} \left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} \lambda + q^{2\nu} \left(2M\mu(1+\mu) - \left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} \lambda \right) - 2q^\nu r^\nu \left[\left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} \lambda + M\mu(\nu-1) \right] \right\} \quad (18)$$

2. Square of Ricci tensor.

The square of the Ricci tensor provides a richer and more detailed measure of spacetime curvature than the Ricci scalar. It captures the interaction between matter-energy distributions and spacetime geometry, offering insights into high-energy astrophysical phenomena, gravitational wave dynamics, and cosmological structure formation. It is also a crucial tool for testing and validating modified gravity theories. Consider the second scalar invariant: the so-called square of Ricci tensor, being responsible for the square of the energy-momentum tensor of a field in the spacetime of a black hole, and it is defined as $\mathcal{R} = R_{\mu\nu} R^{\mu\nu} \equiv 1/(8\pi G) T_{\mu\nu} T^{\mu\nu}$ for the spacetime around black holes with the lapse functions given by Eq. 17. The square of the Ricci tensor takes the following form as

$$\mathcal{R} = \frac{1}{2r^6} \left\{ 5\lambda^2(q^\nu + r^\nu)^4 + 4Mq^\nu \left(1 + \frac{q^\nu}{r^\nu}\right)^{-\frac{2\mu}{\nu}} \times \mu \left[Mq^{3\nu}\mu(5 + (\mu-2)\mu) + \left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} \right] \right\}$$

$$\times (q^\nu + r^\nu)^2 \lambda (q^\nu (\mu - 5) - r^\nu (5 + \nu)) \Big] + \frac{2q^\nu}{r^\nu} ((1 + \nu)(\mu + \nu) - 2\nu) \Big\} \quad (19)$$

3. Kretschmann scalar

The Kretschmann scalar quantifies the total curvature

$$\begin{aligned} \mathcal{K} = \frac{1}{r^6} & \left[13\lambda^2 + \frac{4M\lambda}{\left(1 + \frac{q^\nu}{r^\nu}\right)^{2+\frac{\mu}{\nu}}} \left(10 + \frac{q^{2\nu}}{r^{2\nu}} (\mu - 1)(3\mu - 10) + \frac{q^\nu}{r^\nu} (20 - \mu(13 + 3\nu)) \right) \right. \\ & + 4M^2 \left(1 + \frac{q^\nu}{r^\nu} \right)^{-\frac{2\mu}{\nu}} \left(12 + \frac{q^\nu \mu}{q^\nu + r^\nu} \left(\nu - 20 - \frac{2q^{2\nu}}{(q^\nu + r^\nu)^2} \mu(3 + \nu)(\mu + \nu) \right) \right. \\ & + \frac{q^{3\nu} \mu(\mu + \nu)^2}{(q^\nu + r^\nu)^3} + \frac{q^\nu}{(q^\nu + r^\nu)} (4\nu + \mu(17 + \nu(6 + \nu))) \Big) \Big) \\ & \left. + 4\lambda \ln \frac{r}{|\lambda|} \left(2M \left(1 + \frac{q^\nu}{r^\nu} \right)^{-\frac{\mu}{\nu}} \left(\frac{q^\nu \mu (-q^\nu (\mu - 5) + r^\nu (5 + \nu))}{(q^\nu + r^\nu)^2} - 6 \right) - 5\lambda + 3\lambda \ln \frac{r}{|\lambda|} \right) \right] \quad (20) \end{aligned}$$

Figure 3 illustrates how scalar invariants, such as the Ricci scalar, the squared Ricci tensor, and the Kretschmann scalar, change with distance for various values of q . All invariants decrease rapidly with increasing r and approach the asymptotic value. An increase in the parameter ν causes a slight increase in the Ricci scalar invariant; an increase in the parameter μ causes a slight increase in the Ricci tensor invariant; a decrease in the parameter μ causes a slight increase in the Kretschmann scalar invariant.

III. ENERGY CONDITIONS

Energy conditions are handy tools for discussing cosmological geometry and black hole spacetimes [102–104] in both general relativity [105] and modified gravity [106]. In the models of NED and PFDM, it is essential to consider energy conditions to understand how gravity,

of spacetime and highlights regions of strong-field effects. Near the black hole horizon, the behavior of $\mathcal{K}$ provides insights into the impact of PFDM and NED field parameters on tidal forces. Now we consider the Kretschmann scalar defined as $\mathcal{K} = R_{\mu\nu\sigma\rho} R^{\mu\nu\sigma\rho}$. Usually, the square root of the Kretschmann scalar can be interpreted as an effective gravitational energy density $\sqrt{\mathcal{K}} \sim \rho_M$. One can easily calculate the Kretschmann scalar for the space-time metric in the form

electromagnetic fields, and dark matter behave. Energy conditions are fundamental constraints that ensure matter and energy's physical consistency and stability in theoretical physics and cosmology.

These conditions impose restrictions on the stress-energy tensor, providing a framework to analyze gravitational phenomena without requiring detailed knowledge of the underlying matter dynamics. In the context of NED and PFDM, energy conditions are essential for understanding the interplay between gravity, electromagnetic fields, and dark matter [107].

The four primary energy conditions, WEC, NEC, DEC, and SEC, are particularly significant to study spacetime properties. They ensure non-negative energy densities, causal energy flow, and physically realistic gravitational effects [108]. Additionally, these conditions enable the derivation of significant results, such as singularity theorems, black hole topology theorems, and con-




Fig. 3. (color online) Radial profiles of Scalar invariant for varying values of q . Starting from the left in the top panel, the Ricci scalar, the square of the Ricci tensor, and the Kretschmann scalar, respectively.

straints on cosmic expansion [109].

In NED, energy conditions regulate the behavior of nonlinear electromagnetic fields, preventing unphysical scenarios such as negative energy regions or superluminal propagation. In PFDM, they constrain the properties of dark matter, ensuring its compatibility with observational data and its role in shaping cosmic structures [108]. Together, these frameworks provide a robust foundation for exploring the intricate dynamics of spacetime and the universe's evolution [110].

The standard energy conditions include NEC, WEC, SEC, and DEC, given as

$$\begin{aligned} \text{NEC} : \rho + p_i &\geq 0 \quad (i = r, \theta, \phi); \\ \text{WEC} : \rho &\geq 0 \quad \rho + p_i \geq 0 \quad (i = r, \theta, \phi); \\ \text{SEC} : \rho + \sum_i p_i &\geq 0 \quad \rho + p_i \geq 0 \quad (i = r, \theta, \phi); \\ \text{DEC} : \rho &\geq 0 \quad |p_i| \leq \rho \quad (i = r, \theta, \phi). \end{aligned} \quad (21)$$

Relevant quantities are deduced as

$$\begin{aligned} \rho &= -\frac{2\mu M q^\nu r^\mu}{r^3 (q^\nu + r^\nu)^{\frac{\mu+\nu}{\nu}}} + \frac{\lambda}{8\pi r^3}, \\ \rho + p_r &= 0, \\ \rho + p_{\theta,\phi} &= -\frac{((\mu+1)q^\nu - r^\nu(\nu-1))\mu M q^\nu r^\mu}{r^3 (q^\nu + r^\nu)^{\frac{\mu}{\nu}+2}} + \frac{3\lambda}{16\pi r^3}, \\ \rho + p_r + p_\theta + p_\phi &= \frac{2((\nu+1)r^\nu - (\mu-1)q^\nu)\mu M q^\nu r^\mu}{r^3 (q^\nu + r^\nu)^{\frac{\mu}{\nu}+2}} + \frac{\lambda}{8\pi r^3}, \\ \rho - |p_r| &= 0, \\ \rho - |p_{\theta,\phi}| &= \frac{((\mu-3)q^\nu - (\nu+3)r^\nu)\mu M q^\nu r^\mu}{r^3 (q^\nu + r^\nu)^{\frac{\mu}{\nu}+2}} + \frac{\lambda}{8\pi r^3}. \end{aligned} \quad (22)$$

To examine the energy conditions of NED and PFDM in our case, we plot the unspecified quantities in Eq. (21) concerning the radial coordinate r in Fig. 4.

Fig. 4 shows the different energy states and the dependence of the different energy states on r for different sets of parameters. The graphs depict the distribution of energy density and pressure under the influence of various parameters. As the radius increases, all energy conditions decrease while they attain higher values at smaller distances. An increase in charge reduces energy density, and in some cases, it may become negative, indicating a stronger spacetime curvature due to the electromagnetic field. PFDM expands spacetime and slows down energy distribution changes, but some energy conditions may be violated at high values. The increase in NED parameters affects the internal structure of the black hole, stabilizing energy density and pressure at larger distances. These results are crucial for understanding the distribution of matter, electromagnetic fields, and dark matter around the black hole, revealing the possible existence of exotic or unstable structures in certain cases.

IV. THERMODYNAMICS OF BLACK HOLES SURROUNDED BY PFDM

Solving the equation $f(r_h)=0$, one can obtain the event horizon radius r_h , with which the mass of the black hole can be expressed as

$$M = \frac{1}{2} \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{\frac{\mu}{\nu}} \left(r_h + \lambda \ln \frac{r_h}{|\lambda|} \right) \quad (23)$$

The Hawking temperature T_h is related to the surface gravity by:




Fig. 4. (color online) The variation of ρ , $\rho + p_\theta$, $\rho + p_r + p_\theta + p_\phi$ and $\rho - |p_\theta|$ versus r for NED and PFDM taking $q = 1$, $q=0.2$ and $\lambda = 0.1$.

$$T_h = \frac{f'(r)}{4\pi} \Big|_{r=r_h} = \frac{1}{4\pi} \left[\frac{2M}{r_h^2} \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{-\frac{\mu}{\nu}} + \frac{2M\mu q^\nu}{r_h^{2+\nu}} \right. \\ \left. \times \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{-\frac{\mu}{\nu}-1} - \frac{\lambda}{r_h^2} \left(1 + \ln \frac{r_h}{|\lambda|} \right) \right] \quad (24)$$

In Fig. 5, we plot the temperature of the black hole with NED charge surrounded by PFDM. We can see that the black hole temperature increases and reaches a maximum before having a phase of decrease. Furthermore, this figure shows us that this maximum increases for higher values of the dark matter parameter λ . Let us notice that the case " $\lambda=0$ " corresponds to the temperature of a black hole without dark matter. We write the first law of black hole thermodynamics from the black hole studied here and then find the entropy before computing the corrected entropy. The first law is expressed as

$$dM = T_h dS_0 + \Phi_h dq + \beta_h d\alpha, \quad (25)$$

where S_0 represents the entropy at the equilibrium without considering thermal fluctuation or the uncorrected entropy. Let us notice that S_0 , the magnetic charge q , and the dark matter parameter α form a complete set of extensible variables. T_h is the Hawking temperature at the horizon and Φ_h is the potential. β_h is the conjugating

quantity of the dark matter parameter λ . Now, in the first expression of (25), we can find the formula of the uncorrected entropy S_0 , given by

$$S_0 = \int \frac{1}{T_h} dM = \int \frac{1}{T_h} \frac{\partial M}{\partial r_h} dr_h. \quad (26)$$

To compute this, we have to find $\frac{\partial M}{\partial r_h}$ from Eq.(26). The obtained expression is as follows

$$\frac{\partial M}{\partial r_h} = \frac{1}{2} \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{\frac{\mu}{\nu}} \left(1 + \frac{\lambda}{r_h} \right) - \frac{\mu q^\nu}{2r_h^{\nu+1}} \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{\frac{\mu}{\nu}-1} \left(r_h + \lambda \ln \frac{r_h}{|\lambda|} \right). \quad (27)$$

Introducing Eq.(27) and (24) into Eq. (26), we get the black hole entropy at the equilibrium expressed as

$$S_0 = \frac{2\pi \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{\frac{\mu}{\nu}}}{(q^\nu + r_h^\nu) \left(q^\nu \lambda \left(-1 + \mu \ln \frac{r_h}{|\lambda|} \right) + q^\nu (1 + \mu) r_h - \lambda r_h^\nu + r_h^{1+\nu} \right)^2} \\ \times \left[r_h^{3\nu} (r_h^2 - 2\lambda r_h - \lambda^2) + q^{3\nu} (\lambda^2 (-1 + \mu)) \right]$$




Fig. 5. (color online) The black hole temperature T_h as a function of the event horizon radius r_h , showing the effects of the dark matter parameter λ and the black hole charge q on thermal evolution

$$\begin{aligned}
& -2\lambda(1+\mu^2)r_h + (1+\mu)(-1+\mu)^2r_h^2 - 6\lambda r_h \\
& + (3+\mu(-1+2\nu))r_h^2 + q^{2\nu}\lambda^2\mu^2 \ln \frac{r_h^2}{|\lambda|} (q^\nu(-1+\mu) - r_h^\nu) \\
& + 2q^\nu\lambda\mu \ln \frac{r_h}{|\lambda|} (q^{2\nu}(\lambda - \lambda\mu + (1-\mu+\mu^2)r_h) \\
& + r_h^{2\nu}(\lambda + (1+\nu)r_h) + q^\nu r_h^\nu(-\lambda(-2+\mu) + (2-\mu+\nu)r_h)) \Big] \quad (28)
\end{aligned}$$

Here, we can see that this result is the same as obtained in the presence of quintessence dark energy found in Ref. [111, 112]. As a result, we can say that perfect fluid dark matter does not affect the evolution of the entropy of a non-linear black hole.

Now, we will compute the corrected entropy at the equilibrium S , for which the general formula is expressed as

$$S = S_0 - \beta \ln(S_0 T_h) \quad (29)$$

Here, β is the corrected parameter with only two values. If $\beta = 0$, Eq.(29) describes the uncorrected entropy, and for $\beta = 1/2$, Eq.(29) represents the corrected entropy due to thermal fluctuation.

From the plot Fig. 6, one can compare the corrected and equilibrium entropy for a given black hole. In the absence of the correction parameter β , the equilibrium entropy of the system is a linearly increasing function, as shown in the bottom. For the correction parameter $\beta = 1/2$, the corrected entropy of the system always has a positive value, which is important for the system. The corrected entropy is a decreasing function below the critical horizon radius, but above this point is always an increasing function. Therefore, for a larger black hole, the corrected entropy of the system is not significant as ex-

pected. However, for a smaller black hole system, the correction term significantly affects the system's entropy due to small thermal fluctuations.

A. Specific Heat

The specific heat C at constant volume is given by:

$$C = T_h \left(\frac{\partial S}{\partial T_h} \right) \quad (30)$$

We need the relationship between the entropy S and the Hawking temperature T_h , which involves the implicit dependence on r_h .

We need:

$$\frac{dT_h}{dr_h} = 2\pi r_h \quad (31)$$

Then, we have

$$\begin{aligned}
\frac{dT_h}{dr_h} \Big|_{r_h} &= \frac{1}{4\pi} \left[\frac{d}{dr_h} \left(\frac{2M}{r_h^2} \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{-\frac{\mu}{\nu}} \right) \right. \\
&+ \frac{d}{dr_h} \left(\frac{2M\mu q^\nu}{r_h^{2+\nu}} \left(1 + \frac{q^\nu}{r_h^\nu} \right)^{-\frac{\mu}{\nu}-1} \right) \\
&\left. - \frac{d}{dr_h} \left(\frac{\lambda}{r_h^2} \left(1 + \ln \frac{r_h}{|\lambda|} \right) \right) \right] \quad (32)
\end{aligned}$$

Finally, the specific heat is:

$$C = T_h \frac{dS}{dr_h} \left(\frac{dr_h}{dT_h} \right) \quad (33)$$

After computing it, we get the following expression




Fig. 6. (color online) Variation of entropy (S) with black hole horizon radius (r_+) for different values of q and λ . Here $\beta = 0$ (no correction) is the bottom panel and $\beta = 1/2$ (with correction) is the top panel.

$$C = \frac{-2\pi r_h^2(q^\nu + r_h^\nu) \left(q^\nu \lambda \ln \frac{r_h}{|\lambda|} - 1 \right) + q^\nu(1 + \mu)r_h - \lambda r_h^\nu + r_h^{1+\nu}}{r_h^{2\nu}(r_h - 2\lambda) + q^{2\nu}(r_h - 2\lambda + (r_h + \lambda)\mu) + q^\nu r_h^\nu(\lambda(\mu - 4) + r_h(2 + \mu + \mu\nu)) + q^\nu \lambda \mu (2q^\nu + r_h^\nu(2 + \nu)) \ln \frac{r_h}{|\lambda|}} \quad (34)$$

A black hole undergoes a second-order phase transition in phase transitions that lead the black hole to move from the stable phase ($C > 0$) to the unstable phase ($C < 0$).

Figure 7 presents the variation of the black hole heat capacity C as a function of the horizon radius, demonstrating the thermodynamic stability of the black hole solution under different values of the charge and the PFDM parameter. One can observe from the plots the presence of discontinuities in the heat capacity, which indicate second-order phase transitions. These transitions separate different thermodynamic phases:

The heat capacity is negative for small black holes (r_h below the discontinuity point), implying that they are thermally unstable. This corresponds to an evaporating phase, in which the black hole loses mass via Hawking radiation. The heat capacity for large black holes (r_h above the transition point) becomes positive, signaling a stable equilibrium phase where the black hole can maintain thermal stability.

Increasing the black hole charge shifts the phase transition point, suggesting that stronger NED effects influence the black hole's thermodynamic structure. The in-

crease in charge leads to a more extended stable phase, reducing the likelihood of black hole evaporation. Increasing the PFDM parameter significantly alters the stability properties. A higher value of it shifts the phase transition to a larger event horizon, implying that the presence of PFDM supports stability for larger black holes. This suggests that black holes embedded in PFDM environments are more resistant to evaporation than those in vacuum or standard dark matter models. From an astrophysical perspective, these findings suggest that black holes in environments with significant PFDM influence may have longer lifetimes as the transition to a stable thermodynamic phase occurs at a larger horizon radius. Moreover, PFDM modifies the black hole's thermal response, potentially affecting accretion disk dynamics, black hole mergers, and observational signatures related to thermal fluctuations.

A well-defined second-order phase transition strengthens the argument that such black holes exhibit non-trivial stability properties, distinguishing them from classical Schwarzschild or Reissner-Nordström black holes. Future observational tests, such as those involving black hole shadow measurements or X-ray spectra from




Fig. 7. (color online) Change of the heat capacity C of the non-linear black hole in the background of perfect fluid dark matter

accretion disks, could constrain the parameters q and λ based on their effects on stability and evaporation rates.

V. NEUTRAL PARTICLE MOTION

Neutral particle motion refers to the movement of particles that do not possess a net electric charge. Electric fields do not influence these particles because of their lack of charge. However, they may still be subject to other forces, such as gravitational, magnetic, or particle interactions. Examples of neutral particles include neutrons, neutrinos, and certain atoms or molecules with equal protons and electrons. Understanding the motion of neutral particles is crucial in various scientific fields, including nuclear physics, astrophysics, and materials science. Studying how neutral particles move and interact with their surroundings provides insight into the behavior of matter under different conditions and helps to understand fundamental processes in the universe.

A. Equations of motion

The standard Lagrangian density for the neutral particle with rest mass m moving around a black hole can be written as

$$\mathcal{L}_p = \frac{1}{2} g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu. \quad (35)$$

Due to the existence of the timelike and spacelike Killing vectors, one may introduce the conserved specific energy $\mathcal{E}$ and specific angular momentum $\mathcal{L}$ of the particle in the following form

$$g_{tt} \dot{t} = -\mathcal{E}, \quad g_{\phi\phi} \dot{\phi} = \mathcal{L}. \quad (36)$$

The normalization condition governs the equations of motion for a test particle.

$$g_{\mu\nu} u^\mu u^\nu = \epsilon \quad (37)$$

where ϵ is 0 and -1 for massless and massive particles,

respectively.

For neutral test particles, motion is governed by time-like geodesics of spacetime, and the motion equations can be found using Eq. (37). Considering Eq. (36), one may easily obtain the following equations of motion.

$$\dot{t} = -\frac{\mathcal{E}}{g_{tt}}, \quad (38)$$

$$\dot{r}^2 = \mathcal{E}^2 + g_{rr} \left(1 + \frac{K}{r^2} \right), \quad (39)$$

$$\dot{\theta} = \frac{1}{g_{\theta\theta}} \left(K - \frac{\mathcal{L}^2}{\sin^2 \theta} \right), \quad (40)$$

$$\dot{\phi} = \frac{\mathcal{L}}{g_{\phi\phi}}, \quad (41)$$

where K is the Carter constant associated with the total angular momentum.

Restricting the motion of the particle to the constant plane $\theta = \text{const}$ and $\dot{\theta} = 0$, the Carter constant takes the form $K = \mathcal{L}^2$, and the equation of the radial motion can be expressed in the following standard form

$$\dot{r}^2 = \mathcal{E}^2 - V_{\text{eff}}. \quad (42)$$

where the effective potential of the motion of neutral particles is defined as

$$V_{\text{eff}} = f(r) \left(1 + \frac{\mathcal{L}^2}{r^2 \sin^2 \theta} \right). \quad (43)$$

B. Circular orbits

To study the circular motion of the particles, we use the following conditions:

$$V_{\text{eff}} = \mathcal{E}, \quad V'_{\text{eff}} = 0. \quad (44)$$




Fig. 8. (color online) The effective potential for radial motion of test particles with fixed specific axial angular momentum $\mathcal{L} = \sqrt{20}M$ around a black hole.

To find the values of angular momentum and energy of particles in circular orbits, we used the conditions giv-

en in Eq. (44) to obtain:

$$\mathcal{L}^2 = \frac{r^2 \left(2M(q^\nu - \mu q^\nu + r^\nu) - \lambda(q^\nu + r^\nu) \ln \frac{r}{|\lambda|} \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu} + \lambda(q^\nu + r^\nu) \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu} \right)}{3\lambda(q^\nu + r^\nu) \ln \frac{r}{|\lambda|} \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu} + 2(\mu - 3)Mq^\nu - 6Mr^\nu + (2r - \lambda)(q^\nu + r^\nu) \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu}},$$

$$\mathcal{E}^2 = \frac{2(q^\nu + r^\nu) \left(\left(\lambda \ln \frac{r}{|\lambda|} + r \right) \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu} - 2M \right) \left(\lambda \ln \frac{r}{|\lambda|} - 2M \left(1 + \frac{q^\nu}{r^\nu} \right)^{-\frac{\mu}{\nu}} + r \right)}{r \left(3\lambda(q^\nu + r^\nu) \ln \frac{r}{|\lambda|} \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu} + 2(\mu - 3)Mq^\nu - 6Mr^\nu + (2r - \lambda)(q^\nu + r^\nu) \left(1 + \frac{q^\nu}{r^\nu} \right)^{\mu/\nu} \right)}. \quad (45)$$

Fig. 9 shows radial profiles of energy $\mathcal{E}$ (top panel) and angular momentum $\mathcal{L}$ (middle panel) corresponding to circular orbits. The bottom panel shows the dependence on energy and angular momentum. $\mathcal{E}$ and $\mathcal{L}$ de-

crease with increasing radial distance and then stabilize. As the angular momentum increases, there is a steady increase in energy. As the values of ν and μ increase, the energy $\mathcal{E}$ and the angular momentum $\mathcal{L}$ also increase.




Fig. 9. (color online) Radial dependence of $\mathcal{L}$, $\mathcal{E}$ to vary q with fixed λ (top panel). Plot of $\mathcal{L}$ and $\mathcal{E}$ for λ and q in the lower panel.

C. Stable circular orbits

The stability condition of circular orbits of test particles around a central point of gravitating compact objects is described by the equation $\partial_{rr} V_{\text{eff}} \geq 0$, while the innermost stable circular orbits satisfy $\partial_{rr} V_{\text{eff}} = 0$. This

condition is often applied with the circularity condition in Eq.(44).

The condition $\partial_{rr} V_{\text{eff}} = 0$ leads to the following equation, which is the solution with respect to the radial coordinate r gives innermost stable circular orbit (ISCO) radius.

$$\begin{aligned}
 & 4\left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} (q^\nu + r^\nu)^2 \lambda^2 - 4M^2 \left(1 + \frac{q^\nu}{r^\nu}\right)^{-\frac{\mu}{\nu}} (3r^{2\nu} + q^{2\nu}(\mu-3)(\mu-1) + q^\nu r^\nu(6 + \mu(\nu-4))) \\
 & + 2M (r^{2\nu}(r-4\lambda) - q^{2\nu}(\mu-1)(r-4\lambda + r\mu) + q^\nu r^\nu(4\lambda(\mu-2) + r(2 + \mu\nu))) \\
 & - \lambda \left(\left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} (q^\nu + r^\nu)^2 (r-4\lambda) - 2M (6r^{2\nu} + q^{2\nu}(6 - \mu(4 + \mu)) + q^\nu r^\nu(12 + \mu(\nu-4))) \right) \\
 & \times \ln \frac{r}{|\lambda|} - \frac{3}{2} r^3 \left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} (q^\nu + r^\nu)^3 \lambda^2 \ln^2 \frac{r}{|\lambda|} \left(\left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} (q^\nu + r^\nu) (2r - \lambda) \right. \\
 & \left. + 2M q^\nu (\mu-3) - 6M r^\nu + 3 \left(1 + \frac{q^\nu}{r^\nu}\right)^{\frac{\mu}{\nu}} (q^\nu + r^\nu) \lambda \ln \frac{r}{|\lambda|} \right) = 0
 \end{aligned} \tag{46}$$

The radius of the ISCO radius of the test particles can be calculated numerically by solving Eq.(46). The solution $r = r_{\text{ISCO}}$ has been shown in Fig. 10.

The dependence of the ISCO radius on the parameters q and λ is shown in Fig. 10. As μ and ν increase, r_{ISCO} values become higher. As q increases, r_{ISCO} decreases, and as λ increases, r_{ISCO} first decreases and then increases.

VI. EFFECTIVE METRIC FOR PHOTONS

The electrostatic fields and electromagnetic waves do not interact in linear electrodynamics. In NED, conversely, a photon does not follow null geodesics; instead,

it interacts with the field due to its nonlinearity. The "shifted-null" geodesics in the spacetime of the effective geometry for photons surrounding the black hole, governing the photon motion of black holes in spacetime obtained in GR coupled with NED [49, 113–115]

$$\tilde{g}^{\mu\nu} = g^{\mu\nu} - 4 \frac{\mathcal{L}_{\mathcal{F}\mathcal{F}}}{\mathcal{L}_{\mathcal{F}}} F_\mu^\lambda F_\lambda^\nu, \tag{47}$$

$$\tilde{g}_{\mu\nu} = 16 \frac{\mathcal{L}_{\mathcal{F}\mathcal{F}} F_{\mu\eta} F_\nu^\eta - (\mathcal{L}_{\mathcal{F}} + 2\mathcal{F} \mathcal{L}_{\mathcal{F}\mathcal{F}}) g_{\mu\nu}}{\mathcal{F}^2 \mathcal{L}_{\mathcal{F}\mathcal{F}}^2 - 16(\mathcal{L}_{\mathcal{F}} + \mathcal{F} \mathcal{L}_{\mathcal{F}\mathcal{F}})^2}, \tag{48}$$

in which




Fig. 10. (color online) Dependence of r_{ISCO} on λ for various values of q (left panel) and on q for varying λ (right panel).

$$\mathcal{L}_{\mathcal{F}} = \frac{\partial \mathcal{L}(\mathcal{F})}{\partial \mathcal{F}}, \quad \mathcal{L}_{\mathcal{F}\mathcal{F}} = \frac{\partial^2 \mathcal{L}(\mathcal{F})}{\partial \mathcal{F}^2}. \quad (49)$$

The corresponding photon eikonal equation in the effective geometry associated with a black hole may be stated as

$$\tilde{g}_{\mu\nu} k^\mu k^\nu = 0. \quad (50)$$

In the above expression, k^μ is the four-wave vector related to the photon four-momentum with the relation $p^\mu = \hbar k^\mu$ (in Gaussian units, $\hbar = 1$). In our previous papers [87, 116, 117], by graphical and numerical analyses, we have shown that the difference in shadow sizes, considering the interaction between the photon and the NED field, does not change much. It is smaller than the error in the data of EHT observations of Sgr A* (~14%) and M87* (about 7.14%). Moreover, our primary focus in this work is probing the combined effects of spacetime parameters, NED field charge, and PFDM field on the black hole shadow shape and size.

VII. GEODESIC MOTION AND BLACK HOLE SHADOW

Geodesic motion around a black hole is the most important aspect in forecasting the potential structure of a black hole shadow. Consider a test particle traveling around the black hole with a rest mass of m_0 . Geodesics are easier to find if the constant of motion associated with the symmetry direction is determined. Let y^σ and $v^\sigma = \frac{dx^\sigma}{d\lambda}$ be the vectors along the symmetry's direction and the tangent vector along the curve $x^\sigma = x^\sigma(\tau)$, respectively, where τ is an affine parameter. If the trajectory x^σ is a geodesic [118], we may use the killing vectors to get

$$y^\sigma v_\sigma = \text{constant}. \quad (51)$$

The time-like killing vector for a time-independent metric coefficient is provided by $y^\sigma = (1, 0, 0, 0)$, and the Killing vector for the ϕ direction may be expressed as $y^\sigma = (0, 0, 0, 1)$. By solving equation 51, we can establish the following relationship utilizing the equation as mentioned above as shown below:

$$y^0 v_0 = v^0 = -\mathcal{E}, \quad y^3 v_3 = v_3 = \mathcal{L}. \quad (52)$$

$\mathcal{E}$ and $\mathcal{L}$ indicate a particle's relativistic energy and angular momentum per unit mass. The geodesic equations may now be determined using Equation (52), as illustrated below:

$$u^0 = g^{0\mu} u_\mu = g^{00} u_0 = \frac{\mathcal{E}}{f(r)}, \quad (53)$$

$$u^3 = g^{3\mu} u_\mu = g^{33} u_3 = \frac{\mathcal{L}}{r^2 \sin^2 \theta}. \quad (54)$$

This becomes

$$\frac{dt}{d\tau} = \frac{\mathcal{E}}{f(r)}, \quad \frac{d\phi}{d\tau} = \frac{\mathcal{L}}{r^2 \sin^2 \theta}. \quad (55)$$

The two remaining geodesic equations may be obtained using the Hamilton-Jacobi approach. Consequently, it may be inferred that

$$\frac{\partial S}{\partial \tau} + \frac{1}{2} g^{\mu\sigma} \frac{\partial S}{\partial x^\mu} \frac{\partial S}{\partial x^\sigma} = 0. \quad (56)$$

A proposed approach, such as the one mentioned in reference [119], would result in the following:

$$H = H_r - \mathcal{E}t + \mathcal{L}\phi + H_\theta + \frac{1}{2} m_0^2 \tau. \quad (57)$$

In Eq. (57), H_r and H_θ represent the functions of r and θ , respectively. Introducing the Jacobi action (56) yields the Hamilton-Jacobi equation.

$$H_r = \int^r \frac{\sqrt{X_r}}{r^2 f(r)} dr, \quad H_\theta = \int^\theta \sqrt{X_\theta} d\theta, \quad (58)$$

in which

$$X_r = r^4 \mathcal{E}^2 - r^2 (r^2 m_0^2 + K + \mathcal{L}^2) f(r), \quad (59)$$

$$X_\theta = K - \mathcal{L}^2 \cot^2 \theta. \quad (60)$$

The geodesic equations of motion for a particle in the presence of a non-rotating black hole may be expressed using the relation $H_\theta = \frac{\partial S}{\partial \theta} = \frac{\partial S_\theta}{\partial \theta}$ and $H_r = \frac{\partial S}{\partial r} = \frac{\partial S_r}{\partial r}$.

$$r^2 \frac{d\theta}{d\tau} = \sqrt{K - \mathcal{L}^2 \cot^2 \theta}, \quad (61)$$

$$r^2 \frac{dr}{d\tau} = \sqrt{r^4 \mathcal{E}^2 - r^2 (r^2 m_0^2 + K + \mathcal{L}^2) f(r)}. \quad (62)$$

Eq.(61) represents the geodesic equation along the θ direction, whereas Eq. (62) represents the geodesic equation along the r direction. Furthermore, the subsequent photon-traced geodesic equations are derived by assuming $m_0^2 = 0$ in the vicinity of the black hole [120].

$$r^2 \frac{d\theta}{dr} = \sqrt{K - \mathcal{L}^2 \cot^2 \theta}, \quad (63)$$

$$\beta = \lim_{r \rightarrow \infty} \left(r^2 \frac{d\theta}{dr} \right). \quad (71)$$

$$r^2 \frac{dr}{d\tau} = \sqrt{r^4 \mathcal{E}^2 - r^2 (K + \mathcal{L}^2) f(r)}. \quad (64)$$

On the other hand, we may write Eq. (64) as

$$\left(\frac{dr}{d\tau} \right)^2 + V_{eff} = 0, \quad (65)$$

The symbol V_{eff} represents the effective potential and is given by

$$V_{eff} = \frac{f(r)}{r^2} (K + \mathcal{L}^2) - \mathcal{E}^2. \quad (66)$$

The following limitations will aid us in identifying the unstable circular orbits:

$$V_{eff}(r = r_p) = V'_{eff}(r = r_p) = 0, \quad (67)$$

In this context, $r = r_p$ denotes the radius of a photon, and ' indicates the derivative concerning r . Hence, the equation $V_{eff}(r_p) = 0$ indicates that

$$\frac{r_p^2}{f(r_p)} = \xi + \eta^2. \quad (68)$$

Equation (68) make use of the Chandrasekhar constant definition [119], which is $\xi = K/\mathcal{E}^2$ and $\eta = \mathcal{L}/\mathcal{E}$. Similarly, $V'_{eff}(r_p) = 0$, results in

$$r_p f'(r_p) - 2f(r_p) = 0. \quad (69)$$

Imagine a massless particle, like a photon, released by an object traveling toward the black hole between an observer and a brilliant object. Photon paths may result in (a) scattering away, (b) falling into the black hole, or (c) a critical geodesic dividing (a) and (b). The scattering of a photon from the black hole is visible to the viewer during this entire process. However, a dark zone will develop when a photon enters the black hole. We call this dark zone the shadow of the black hole. Our primary objective in this part is to investigate the shadow emitted in PFDM spacetime by the black hole. Initially, we must specify the celestial coordinates [121–123] for this purpose as

$$\alpha = \lim_{r \rightarrow \infty} - \left(r^2 \sin \theta \frac{d\phi}{dr} \right), \quad (70)$$

In this context, α represents the apparent perpendicular distance of the shadow from the axis of symmetry, whereas β represents its projection on the equatorial plane. The variable θ denotes the inclination angle between the axis of symmetry and the observer's line of sight, while r_0 indicates the distance between the black hole and the observer. The values of the derivatives $d\phi/dr$ and $d\theta/dr$ may be determined by using the geodesic equations in the following manners:

$$\frac{d\phi}{dr} = \frac{\mathcal{L} \csc^2 \theta}{r^2 \sqrt{\mathcal{E}^2 - \frac{f(r)}{r^2} (K + \mathcal{L}^2)}}, \quad (72)$$

$$\frac{d\theta}{dr} = \frac{1}{r^2} \sqrt{\frac{K - \mathcal{L}^2 \cot^2 \theta}{\mathcal{E}^2 - \frac{f(r)}{r^2} (K + \mathcal{L}^2)}}. \quad (73)$$

When Eqs. (72) and (73) are solved for $r \rightarrow \infty$, the expression for the celestial coordinates (70) assumes the following form [124, 125]:

$$\alpha = -\eta \csc^2(\theta) \sin(\theta), \quad \beta = \pm \sqrt{\xi - \eta^2 \cot^2(\theta)}. \quad (74)$$

The preceding equations are further simplified when $\theta = \pi/2$ is substituted in.

$$\alpha = -\eta, \quad \beta = \pm \sqrt{\xi}. \quad (75)$$

As a result, the expression of the shadow radius in the celestial plane (α, β) may be computed as:

$$R_s^2 = \alpha^2 + \beta^2 = \eta^2 + \xi = \frac{r_p^2}{f(r_p)} \quad (76)$$

In the equation (76), R_s stands for the radius of the black hole's shadow while it is not revolving.

VIII. CONSTRAINTS ON NED AND PFDM PARAMETERS OF THE BLACK HOLE WITH THE EHT OBSERVATIONS OF M87 AND Sgr. A

In this section, we examine the dependence of the PFDM parameter λ and q on the shadow images of the black holes M87* and Sagittarius A* (Sgr. A*) using Event Horizon Telescope (EHT) observations. Our analysis is




Fig. 11. (color online) The radial profile of black hole's shadow for parameter q (left panel), parameter λ (middle panel) and M (right panel) at different combinations of parameters μ and $\nu = 1$.

limited to the non-rotating case, since the rotation parameter a of Sgr. A* is small enough to have a negligible effect on the shadow radius [126]. Furthermore, it is shown that it is difficult to distinguish between a Kerr

black hole ($a = 0.60M$) and a non-rotating extended black hole for M87* using shadow images alone. This conclusion is supported by general-relativistic magnetohydrodynamic simulations and radiative transfer calculations




Fig. 12. (color online) Dependence of r_{ph} on λ for various values of q (top panel) and on q for varying λ (bottom panel).

that generate synthetic shadow images for direct comparison with EHT observations [127].

Furthermore, as noted in Ref. [128], the shadow size of M87* is observed to lie in the range of $3\sqrt{3}(1 \pm 0.17)M$.

Regardless of whether the underlying model assumes spherical symmetry or axisymmetric. This observation provides a strong basis for constraining theoretical models of black holes in alternative theories of general relativity and gravity. By comparing theoretical predictions of the shadow radius from our model with EHT observational data, we attempt to determine the allowable ranges for the parameter q and other relevant physical parameters. This approach allows us to probe the interplay between nonlinear electrodynamics, dark matter, and gravitational effects around black holes in the strong field regime.

The EHT collaboration reported the following data for M87*: the angular diameter of the shadow is $\theta_{M87*} = 42 \pm 3 \mu\text{as}$, the distance to M87* is $D = 16.8 \text{ Mpc}$, and the mass of M87* is $M_{M87*} = 6.5 \pm 0.90 \times 10^9 M_{\odot}$ [129]. For Sgr. A*, recent EHT results provide the angular diameter of the shadow as $\theta_{\text{Sgr. A}^*} = 48.7 \pm 7 \mu\text{as}$, the distance to Sgr. A* as $D = 8277 \pm 33 \text{ pc}$, and the black hole mass as $M_{\text{Sgr. A}^*} = 4.3 \pm 0.013 \times 10^6 M_{\odot}$ (VLTI) [130, 131].

Using the provided data, the shadow diameter in units of mass can be calculated using the formula [132]:

$$d_{\text{sh}} = \frac{D\theta}{M}. \quad (77)$$

The theoretical shadow diameter is given by $d_{\text{sh}} = 2R_{\text{sh}}$. Applying the above expression, we find the shadow diameter for M87* to be $d_{\text{sh}}^{M87*} = (11 \pm 1.5)M$ and

for Sgr. A* to be $d_{\text{sh}}^{\text{Sgr. A}^*} = (9.5 \pm 1.4)M$.

The relationship between q and λ affecting the black hole shadow size is shown in Fig. 13. The graphs are drawn based on EHT observational data for M87* and Sgr. A*, with the q/M axis representing the charge-to-mass ratio and the d_{sh}/M axis representing the shadow diameter-to-mass ratio. The black lines correspond to $\lambda = 0$, representing the scenario where the effect of dark matter is not considered. The red dashed lines show the variation in shadow size when $\lambda = 0.03$, illustrating how the presence of dark matter influences the shadow. It is visible that as q/M increases, the shadow diameter decreases and this reduction occurs more rapidly when λ is present. The confidence intervals 1σ and 2σ represent the observational data provided by the EHT. When λ is absent, the black lines mostly fall within the 1σ confidence interval, indicating that the model without the influence of dark matter aligns well with the EHT observations. In contrast, the red dashed lines representing the case where λ is present mostly fall within the 2σ interval, suggesting the possible influence of dark matter. However, this influence is not as strong within the 1σ range. Thus, the graphs clearly show how the parameters q and λ affect the black hole shadow size. An increase in charge and the dark matter parameter reduces the shadow size. Based on EHT observations, the absence of λ corresponds more closely to the 1σ interval, indicating a minimal influence of dark matter. However, the presence of λ aligns with the 2σ interval, suggesting the possible existence of dark matter.

IX. CONCLUSION

This study investigated the properties of black holes in GR coupled to NED in the presence of PFDM. The main results obtained across different sections are sum-




Fig. 13. (color online) The plot shows the constraints for q . The corresponding values of q at the mean, 1σ , and 2σ confidence levels are shown.

marized below: A singular black hole solution was derived, incorporating the effects of NED and PFDM. The energy conditions, including the WEC, NEC, SEC, and DEC, are examined. It was observed that as the radius increases, all energy conditions decrease. Higher charge

values reduce energy density and sometimes lead to negative energy densities, indicating the influence of strong spacetime curvature. The PFDM parameter increase slows energy distribution changes, whereas NED parameters stabilize energy density at more considerable dis-

tances.

The thermodynamic properties, including temperature, entropy, and specific heat, are analyzed. The specific heat capacity transitioned from positive to negative values, suggesting stability and phase transition behavior in particular parameter ranges. PFDM and NED modified the black hole's Hawking temperature, indicating changes in its thermodynamic equilibrium.

The geodesic motion of neutral particles was studied, focusing on circular orbits. ISCO was derived from the black hole's parameters. It was observed that increasing PFDM parameters shifts the ISCO radius, while higher charge values decrease it.

The black hole shadow was computed under different parameter values, revealing how PFDM and NED influence its size and shape. The radius of the shadow increased with higher values of PFDM parameters but decreased with increasing charge. The computed results provide insights into how black hole imaging techniques can place future observational constraints on these parameters.

The results demonstrated that the influence of NED and PFDM on black hole properties: Show the dependence of energy conditions on charge and PFDM parameters. Notable deviations from standard GR expectations were identified. Depict the variations in temperature, entropy, and specific heat, highlighting regions of stability and instability. Present the shifts in circular orbits and ISCO under different parameter values, showing a direct influence of PFDM and NED. We have also demonstrated how to modify observational characteristics such as the shadow radius, which is crucial for astrophysical applications.

The study provides a new framework for understanding black holes in alternative gravity models with dark matter interactions. The findings suggest that black holes in PFDM environments exhibit modified thermodynamic and observational properties. Future black hole imaging and gravitational wave observations can test these properties. Further research is needed to explore the role of rotation and higher-order corrections in these models. The results can be extended to analyze accretion disk properties and their implications for high-energy astrophysics. In conclusion, this work contributes to understanding black holes in modified gravity scenarios and offers potential pathways for observational tests using astrophysical and cosmological data.

Our analysis of the black hole shadow in the presence of NED and PFDM has provided key insights into how these parameters affect the observable shadow radius. Comparing our results with the EHT measurements of M87* and Sgr A*, we find that PFDM contributes to an increase in the shadow size. At the same time, NED effects introduce modifications that can reduce it. The interplay of these effects allows us to place constraints on the model parameters, ensuring that deviations from General Relativity remain within observational limits. Precisely, we determine that the values of NED charge parameter q and PFDM density parameter λ must be finely tuned to reproduce the observed shadow sizes within the 7.14% (M87) and 14% (Sgr A) uncertainty ranges reported by EHT. These constraints provide a novel way to test alternative gravity models and highlight the potential detectability of PFDM signatures in future high-precision black hole imaging experiments.

References

- [1] A. D. Sakharov, Soviet Journal of Experimental and Theoretical Physics **22**, 241 (1966)
- [2] E. B. Gliner, Soviet Journal of Experimental and Theoretical Physics **22**, 378 (1966)
- [3] L. E. Gurevich, Astrophysics and Space Science **38**, 67 (1975)
- [4] A. A. Starobinskiĭ, Soviet Journal of Experimental and Theoretical Physics Letters **30**, 682 (1979)
- [5] Y. Kurmanov, K. Boshkayev, T. Konysbayev, O. Luongo, N. Saiyp, A. Urzazalina, G. Ikhsan, and G. Suliyeva, Physics of the Dark Universe **46**, 101566 (2024), arXiv: [2404.15437\[gr-qc\]](#)
- [6] C. Lämmerzahl, M. Maceda, and A. Macías, Classical and Quantum Gravity **36**, 015001 (2019), arXiv: [1802.03766\[gr-qc\]](#)
- [7] S. G. Ghosh and R. K. Walia, Annals of Physics **434**, 168619 (2021), arXiv: [2109.13031\[gr-qc\]](#)
- [8] Planck Collaboration, Astronomy and Astrophysics **641**, A6 (2020), arXiv: [1807.06209\[astro-ph.CO\]](#)
- [9] A. Das, A. Saha, and S. Gangopadhyay, Classical and Quantum Gravity **38**, 065015 (2021)
- [10] V. V. Kiselev, Classical and Quantum Gravity **20**, 1187 (2003), arXiv: [gr-qc/0210040\[gr-qc\]](#)
- [11] B. Toshmatov, Z. Stuchlík, and B. Ahmedov, European Physical Journal Plus **132**, 98 (2017)
- [12] F. Rahaman, K. K. Nandi, A. Bhadra, M. Kalam, and K. Chakraborty, Physics Letters B **694**, 10 (2010), arXiv: [1009.3572\[gr-qc\]](#)
- [13] Y. Zhang and Y. X. Gui, Classical and Quantum Gravity **23**, 6141 (2006), arXiv: [gr-qc/0612009\[gr-qc\]](#)
- [14] S. G. Ghosh, S. D. Maharaj, D. Baboolal, and T.-H. Lee, European Physical Journal C **78**, 90 (2018), arXiv: [1708.03884\[gr-qc\]](#)
- [15] S. U. Khan and J. Ren, Physics of the Dark Universe **30**, 100644 (2020)
- [16] C. A. Benavides-Gallego, A. Abdujabbarov, and C. Bambi, Phys. Rev. D **101**, 044038 (2020)
- [17] B. Narzilloev, J. Rayimbaev, S. Shaymatov, A. Abdujabbarov, B. Ahmedov, and C. Bambi, Phys. Rev. D **102**, 104062 (2020), arXiv: [2011.06148\[gr-qc\]](#)
- [18] S. U. Khan and J. Ren, Chinese Journal of Physics **78**, 141 (2022)
- [19] Mustafa, G. and Hussain, Ibrar, European Physical Journal C **81**, 419 (2021)

- [20] Mustafa, G. and Atamurotov, Farruh and Hussain, Ibrar and Shaymatov, Sanjar and Övgün, A. li, *Chinese Physics C* **46**, 125107 (2022), arXiv: [2207.07608\[gr-qc\]](#)
- [21] M. Zahid, J. Rayimbaev, F. Sarikulov, S. U. Khan, and J. Ren, *European Physical Journal C* **83**, 855 (2023)
- [22] S. U. Khan, J. Rayimbaev, F. Sarikulov, and O. Abdurakhmonov, *Chinese Journal of Physics* **90**, 690 (2024), arXiv: [2310.05860\[gr-qc\]](#)
- [23] M.-H. Li and K.-C. Yang, *Phys. Rev. D* **86**, 123015 (2012), arXiv: [1204.3178\[astro-ph.CO\]](#)
- [24] X. Hou, Z. Xu, and J. Wang, *Journal of Cosmology and Astroparticle Physics* **2018**, 040 (2018), arXiv: [1810.06381\[gr-qc\]](#)
- [25] S. U. Khan, M. Shahzadi, and J. Ren, *Physics of the Dark Universe* **26**, 100331 (2019), arXiv: [2005.09415\[gr-qc\]](#)
- [26] R. A. Konoplya, *Physics Letters B* **795**, 1 (2019), arXiv: [1905.00064\[gr-qc\]](#)
- [27] S. Haroon, M. Jamil, K. Jusufi, K. Lin, and R. B. Mann, *Phys. Rev. D* **99**, 044015 (2019), arXiv: [1810.04103\[gr-qc\]](#)
- [28] M. Akramov, B. Eshchanov, S. Usanov, S. Norbekov, and D. Matrasulov, *Physics Letters A* **524**, 129827 (2024)
- [29] K. Jusufi, M. Jamil, and T. Zhu, *European Physical Journal C* **80**, 354 (2020), arXiv: [2005.05299\[gr-qc\]](#)
- [30] J. Rayimbaev, S. Shaymatov, and M. Jamil, *European Physical Journal C* **81**, 699 (2021), arXiv: [2107.13436\[gr-qc\]](#)
- [31] A. Kumar, A. Sood, J. K. Singh, A. Beesham, and S. G. Ghosh, *Physics of the Dark Universe* **40**, 101220 (2023)
- [32] G. Rakhimova, F. Atamurotov, F. Javed, A. Abdujabbarov, and G. Mustafa, *Nuclear Physics B* **996**, 116363 (2023)
- [33] Mustafa, G. and Javed, Faisal and Maurya, S. K. and Ray, Saibal, *Chinese Journal of Physics* **88**, 32 (2024), arXiv: [2211.10778\[gr-qc\]](#)
- [34] J. M. Bardeen, B. Carter, and S. W. Hawking, *Communications in Mathematical Physics* **31**, 161 (1973)
- [35] S. W. Hawking, *Communications in Mathematical Physics* **43**, 199 (1975)
- [36] S. W. Hawking, *Phys. Rev. D* **13**, 191 (1976)
- [37] D. Kubizňák, R. B. Mann, and M. Teo, *Classical and Quantum Gravity* **34**, 063001 (2017), arXiv: [1608.06147\[hep-th\]](#)
- [38] I. Kaygorodov, A. Khudoyberdiyev, and Z. Shermatova, *Journal of Geometry and Physics* **207**, 105356 (2025), arXiv: [2406.15532\[math.RA\]](#)
- [39] B. Narzilloev and B. Ahmedov, **98**, 101922 (2023).
- [40] M. Rizwan and K. Jusufi, *European Physical Journal C* **83**, 944 (2023), arXiv: [2310.15182\[gr-qc\]](#)
- [41] A. A. Abdujabbarov, B. J. Ahmedov, and V. G. Kagramanova, *General Relativity and Gravitation* **40**, 2515 (2008), arXiv: [0802.4349\[gr-qc\]](#)
- [42] V. P. Frolov, *Phys. Rev. D* **85**, 024020 (2012), arXiv: [1110.6274\[gr-qc\]](#)
- [43] J. Kovář, P. Slaný, C. Cremaschini, Z. Stuchlík, V. Karas, and A. Trova, *Phys. Rev. D* **90**, 044029 (2014), arXiv: [1409.0418\[gr-qc\]](#)
- [44] A. Tursunov, Z. Stuchlík, and M. Kološ, *Phys. Rev. D* **93**, 084012 (2016), arXiv: [1603.07264\[gr-qc\]](#)
- [45] Z. Stuchlík and M. Kološ, *European Physical Journal C* **76**, 32 (2016), arXiv: [1511.02936\[gr-qc\]](#)
- [46] M. Akramov, C. Trunk, J. Yusupov, and D. Matrasulov, *EPL (Europhysics Letters)* **147**, 62001 (2024), arXiv: [2411.14397\[quant-ph\]](#)
- [47] J. Vrba, A. Abdujabbarov, A. Tursunov, B. Ahmedov, and Z. Stuchlík, *European Physical Journal C* **79**, 778 (2019), arXiv: [1909.12026\[gr-qc\]](#)
- [48] J. Rayimbaev, M. Figueroa, Z. Stuchlík, and B. Juraev, *Phys. Rev. D* **101**, 104045 (2020)
- [49] S. U. Khan and J. Ren, *Chinese Journal of Physics* **70**, 55 (2021), arXiv: [2010.02754\[gr-qc\]](#)
- [50] J. Rayimbaev, S. Shaymatov, and M. Jamil, *European Physical Journal C* **81**, 699 (2021), arXiv: [2107.13436\[gr-qc\]](#)
- [51] M. Zahid, S. U. Khan, J. Ren, and J. Rayimbaev, *International Journal of Modern Physics D* **31**, 2250058 (2022)
- [52] S. U. Khan and Z.-M. Chen, *European Physical Journal C* **83**, 704 (2023)
- [53] B. Turimov, S. Usanov, and Y. Khamroev, arXiv eprints, arXiv: 2502.11185 (2025), arXiv: [2502.11185\[gr-qc\]](#).
- [54] S. U. Khan, J. Ren, and J. Rayimbaev, *Modern Physics Letters A* **37**, 2250064 (2022)
- [55] S. U. Khan, U. Uktamov, J. Rayimbaev, A. Abdujabbarov, I. Ibragimov, and Z.-M. Chen, *European Physical Journal C* **84**, 203 (2024)
- [56] S. U. Khan, J. Rayimbaev, F. Sarikulov, and O. Abdurakhmonov, *Chinese Journal of Physics* **90**, 690 (2024), arXiv: [2310.05860\[gr-qc\]](#)
- [57] B. Toshmatov, O. Rahimov, B. Ahmedov, and A. Ahmedov, *Galaxies* **9**, 65 (2021)
- [58] B. Ahmedov, O. Rahimov, and B. Toshmatov, *Universe* **7**, 307 (2021)
- [59] O. Rahimov, B. Toshmatov, Y. Vybylyi, A. Akhmedov, and B. Abdulazizov, *Physics of the Dark Universe* **44**, 101483 (2024)
- [60] B. Turimov and O. Rahimov, *Universe* **8**, 507 (2022)
- [61] B. Turimov, O. Rahimov, B. Ahmedov, Z. Stuchlík, and K. Boymurodova, *International Journal of Modern Physics D* **30**, 2150037 (2021)
- [62] B. Turimov, A. Mamadjanov, and O. Rahimov, *Galaxies* **11**, 70 (2023)
- [63] B. Rahmatov, M. Zahid, J. Rayimbaev, R. Rahim, and S. Murodov, *Chin. J. Phys.* **92**, 143 (2024)
- [64] Z. Hu, Z. Zhong, P.-C. Li, M. Guo, and B. Chen, *Phys. Rev. D* **103**, 044057 (2021), arXiv: [2012.07022\[gr-qc\]](#)
- [65] Z. Zhong, Z. Hu, H. Yan, M. Guo, and B. Chen, *Phys. Rev. D* **104**, 104028 (2021), arXiv: [2108.06140\[gr-qc\]](#)
- [66] S.-b. Yuan, C.-k. Luo, Z.-z. Hu, Z.-y. Zhang, and B. Chen, *Chinese Physics C* **49**, 025103 (2025), arXiv: [2403.06886\[gr-qc\]](#)
- [67] M. Zhang and M. Guo, *European Physical Journal C* **80**, 790 (2020), arXiv: [1909.07033\[gr-qc\]](#)
- [68] J. L. Synge, *Mon. Not. Roy. Astron. Soc.* **131**, 463 (1966)
- [69] J. P. Luminet, *Astronomy and Astrophysics* **75**, 228 (1979)
- [70] J. M. Bardeen, *Proceedings, Ecole d'Été de Physique Théorique: Les Astres Occlus : Les Houches, France, August, 1972*, 215-240, 215 (1973).
- [71] S. Chandrasekhar, *The mathematical theory of black holes* (1983).
- [72] A. de Vries, *Classical and Quantum Gravity* **17**, 123 (2000)
- [73] K. Hioki and K.-i. Maeda, *Phys. Rev. D* **80**, 024042 (2009)
- [74] A. A. Abdujabbarov, L. Rezzolla, and B. J. Ahmedov, *Monthly Notices of the Royal Astronomical Society* **454**, 2423 (2015), arXiv: [1503.09054\[gr-qc\]](#)
- [75] R. Takahashi, *Publications of the Astronomical Society of Japan* **57**, 273 (2005), arXiv: [astro-ph/0505316\[astro-ph\]](#)

- [76] S.-W. Wei and Y.-X. Liu, *Journal of Cosmology and Astroparticle Physics* **2013**, 063 (2013), arXiv: [1311.4251\[gr-qc\]](#)
- [77] A. Abdujabbarov, F. Atamurotov, Y. Kucukakca, B. Ahmedov, and U. Camci, *Astrophysics and Space Science* **344**, 429 (2013), arXiv: [1212.4949\[physics.gen-ph\]](#)
- [78] L. Amarilla and E. F. Eiroa, *Phys. Rev. D* **85**, 064019 (2012)
- [79] L. Amarilla and E. F. Eiroa, *Phys. Rev. D* **87**, 044057 (2013)
- [80] F. Atamurotov, A. Abdujabbarov, and B. Ahmedov, *Phys. Rev. D* **88**, 064004 (2013)
- [81] A. Grenzebach, V. Perlick, and C. Lämmerzahl, *Phys. Rev. D* **89**, 124004 (2014), arXiv: [1403.5234\[gr-qc\]](#)
- [82] U. Papnoi, F. Atamurotov, S. G. Ghosh, and B. Ahmedov, *Phys. Rev. D* **90**, 024073 (2014), arXiv: [1407.0834\[gr-qc\]](#)
- [83] P. V. P. Cunha, C. A. R. Herdeiro, E. Radu, and H. F. Runarsson, *Phys. Rev. Lett.* **115**, 211102 (2015), arXiv: [1509.00021\[gr-qc\]](#)
- [84] Event Horizon Telescope Collaboration, *Astrophysical Journal Letters* **875**, L5 (2019), arXiv: [1906.11242\[astro-ph.GA\]](#)
- [85] The Event Horizon Telescope Collaboration, arXiv e-prints, arXiv: [2311.08680](#) (2023), arXiv: [2311.08680\[astro-ph.HE\]](#)
- [86] M. A. Raza, J. Rayimbaev, F. Sarikulov, M. Zubair, B. Ahmedov, and Z. Stuchlík, *Physics of the Dark Universe* **44**, 101488 (2024), arXiv: [2311.15784\[gr-qc\]](#)
- [87] A. Abdujabbarov, B. Toshmatov, Z. Stuchlík, and B. Ahmedov, arXiv e-prints, arXiv: [1512.05206](#) (2015), arXiv: [1512.05206\[gr-qc\]](#)
- [88] C. Promsiri, W. Horinouchi, and E. Hirunsirisawat, arXiv e-prints, arXiv: [2409.01582](#) (2024), arXiv: [2409.01582\[gr-qc\]](#)
- [89] S. Guo, G.-R. Li, and G.-P. Li, *Chinese Physics C* **46**, 095101 (2022), arXiv: [2205.04957\[gr-qc\]](#)
- [90] V. Vertogradov and A. Övgün, *Classical and Quantum Gravity* **42**, 025024 (2024)
- [91] K. A. Bronnikov, *Physical Review D* **63**, 044005 (2001)
- [92] C. Bambi, D. Malafarina, and L. Modesto, *Physical Review D* **88** (2013), 10.1103/physrevd.88.044009.
- [93] E. Ayón-Beato and A. García, *Physical Review Letters* **80**, 5056 (1998)
- [94] E. Ayón-Beato, *Physics Letters B* **464**, 25 (1999)
- [95] E. Ayón-Beato and A. García, *General Relativity and Gravitation* **31**, 629 (1999)
- [96] C. Bambi and L. Modesto, *Physics Letters B* **721**, 329 (2013)
- [97] J. Rayimbaev, M. Figueroa, Z. c. v. Stuchlík, and B. Juraev, *Phys. Rev. D* **101**, 104045 (2020)
- [98] Z.-Y. Fan and X. Wang, *Physical Review D* **94**, 124027 (2016)
- [99] J. S. B. Toshmatov, Z. Stuchlík and B. Ahmedov, *Physical Review D* **97**, 084058 (2018)
- [100] Z. S. B. Toshmatov and B. Ahmedov, *Physical Review D* **98**, 085021 (2018)
- [101] Z.-Y. Fan and X. Wang, *Physical Review D* **94** (2016), 10.1103/physrevd.94.124027.
- [102] O. Zaslavskii, *Physics Letters B* **688**, 278 (2010)
- [103] J. Neves and A. Saa, *Physics Letters B* **734**, 44 (2014)
- [104] S. Rakhmanov, K. Matchonov, H. Yusupov, K. Nasriddinov, and D. Matrasulov, *The European Physical Journal B* **98**, 35 (2025)
- [105] E.-A. Kontou and K. Sanders, *Classical and Quantum Gravity* **37**, 193001 (2020)
- [106] S. Capozziello, F. S. Lobo, and J. P. Mimoso, *Physics Letters B* **730**, 280 (2014)
- [107] M. Novello, S. E. Perez Bergliaffa, and J. Salim, *Physical Review D* **69** (2004), 10.1103/physrevd.69.127301.
- [108] J. M. Bardeen, B. Carter, and S. W. Hawking, *Commun. Math. Phys.* **31**, 161 (1973)
- [109] R. Penrose, *Phys. Rev. Lett.* **14**, 57 (1965)
- [110] V. SAHNI and A. STAROBINSKY, *International Journal of Modern Physics D* **09**, 373 (2000)
- [111] R. B. T. Ndongmo, S. Mahamat, T. B. Bouetou, C. B. Tabi, and T. C. Kofane, *International Journal of Theoretical Physics* **63** (2024), 10.1007/s10773-024-05687-8.
- [112] B. Turimov, S. Usanov, and Y. Khamroev, *Physics of the Dark Universe* **48**, 101876 (2025), arXiv: [2502.11185\[gr-qc\]](#)
- [113] I. Dymnikova, *Classical and Quantum Gravity* **21**, 4417 (2004), arXiv: [gr-qc/0407072\[gr-qc\]](#)
- [114] Z. Stuchlík and J. Schee, *European Physical Journal C* **79**, 44 (2019)
- [115] J. Schee and Z. Stuchlík, *The Astrophysical Journal* **874**, 12 (2019)
- [116] M. Zahid, F. Sarikulov, C. Shen, M. Umaraliyev, and J. Rayimbaev, *Physics of the Dark Universe* **46**, 101616 (2024)
- [117] M. Zubair, M. A. Raza, F. Sarikulov, and J. Rayimbaev, *Journal of Cosmology and Astroparticle Physics* **2023**, 058 (2023), arXiv: [2305.16888\[gr-qc\]](#)
- [118] D. J. Raine and E. G. Thomas, *Black holes: an introduction* (Imperial College Press, 2010).
- [119] S. Chandrasekhar, *The mathematical theory of black holes*, Vol. 69 (Oxford university press, 1998).
- [120] S.-W. Wei and Y.-X. Liu, *The European Physical Journal Plus* **136**, 436 (2021)
- [121] U. Papnoi, F. Atamurotov, S. G. Ghosh, and B. Ahmedov, *Physical Review D* **90**, 024073 (2014)
- [122] M. Zahid, S. U. Khan, and J. Ren, *Chinese Journal of Physics* **72**, 575 (2021), arXiv: [2101.07673\[gr-qc\]](#)
- [123] M. Zahid, F. Sarikulov, C. Shen, M. Umaraliyev, and J. Rayimbaev, *Phys. Dark Univ.* **46**, 101616 (2024)
- [124] M. Zahid, F. Sarikulov, C. Shen, J. Rayimbaev, K. Badalov, and S. Muminov, *Chin. J. Phys.* **91**, 45 (2024)
- [125] M. Zahid, F. Sarikolov, S. Chao, J. Rayimbaev, and S. Ahmedov, *Classical and Quantum Gravity* (2024).
- [126] S. Vagnozzi, R. Roy, Y.-D. Tsai, and L. Visinelli, (2022), arXiv: [2205.07787\[gr-qc\]](#).
- [127] Y. Mizuno, Z. Younsi, C. M. Fromm, O. Porth, M. De Laurentis, H. Olivares, H. Falcke, M. Kramer, and L. Rezzolla, *Nature Astron.* **2**, 585 (2018), arXiv: [1804.05812\[astro-ph.GA\]](#)
- [128] P. Kocherlakota *et al.* (Event Horizon Telescope), *Phys. Rev. D* **103**, 104047 (2021), arXiv: [2105.09343\[gr-qc\]](#)
- [129] K. Akiyama *et al.* (Event Horizon Telescope), *Astrophys. J. Lett.* **875**, L1 (2019), arXiv: [1906.11238\[astro-ph.GA\]](#)
- [130] D. Wang, (2022), arXiv: [2205.08026\[gr-qc\]](#).
- [131] K. Akiyama *et al.* (Event Horizon Telescope), *Astrophys. J. Lett.* **930**, L12 (2022)
- [132] C. Bambi, K. Freese, S. Vagnozzi, and L. Visinelli, *Phys. Rev. D* **100**, 044057 (2019), arXiv: [1904.12983\[gr-qc\]](#)